

On Double Integrals

V.B.L.Chaurasia, J.C. Arya

-
1. Department of Mathematics, University of Rajasthan, Jaipur-302004 (Rajasthan) India, E-mail: drvble@yahoo.com
 2. Department of Mathematics, Govt. Post-Graduate College, Neemuch (M.P.) India, E-mail: profarya76@gmail.com

ABSTRACT

The present paper is related with the study of the integration of certain products of \bar{H} -function [7], M-series [9] and a general class of polynomials [11]. During the course of finding, we establish certain new double integration involving to a product of a general class of polynomial, M-series and \bar{H} -functions. The results of this paper are believed to be new and basic in nature and are likely to find useful application in various fields notably electrical networks, probability theory and statistical mechanics etc.

Key Words: Feynman integrals, \bar{H} -function, M-series, A general class of polynomials.

2000 Mathematics Subject Classification: 33C60, 26A33.

1. INTRODUCTION

The \bar{H} -function [7] is a generalization of the well known Fox's H-function [4]. The \bar{H} -function pertains to exact partition function of the Gaussian model in statistical mechanics, functions useful in testing hypothesis and several other as its special cases.

The \bar{H} -function will be defined as [1]:

$$\bar{H}_{p,q}^{m,n}[z] = \bar{H}_{p,q}^{m,n} \left[z \left| \begin{matrix} (e_j, \gamma_j; E_j)_{1,n}, (e_j, \gamma_j)_{n+1,p} \\ (f_j, \delta_j)_{1,m}, (f_j, \delta_j; F_j)_{m+1,q} \end{matrix} \right. \right]$$

$$= \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \bar{\phi}(\xi) z^\xi d\xi \quad \dots(1.1)$$

where

$$\bar{\phi}(\xi) = \frac{\prod_{j=1}^m \Gamma(f_j - \delta_j \xi) \prod_{j=1}^n \{\Gamma(1 - e_j + \gamma_j \xi)\}^{E_j}}{\prod_{j=m+1}^q \{\Gamma(1 - f_j + \delta_j \xi)\}^{F_j} \prod_{j=n+1}^p \Gamma(e_j - \gamma_j \xi)} \quad \dots(1.2)$$

which contains fractional powers of some of the gamma functions. Here and throughout the paper e_j ($j = 1, \dots, p$) and f_j ($j = 1, \dots, q$) are complex parameters, $\gamma_j \geq 0$ ($j = 1, \dots, q$), δ_j ($j = 1, \dots, q$) (not all zero simultaneously) and the exponents E_j ($j = 1, \dots, n$) and F_j ($j = m+1, \dots, q$) can take non-integer values.

The \bar{H} -function in (1) converges absolutely if

$$|\arg(z)| < \frac{1}{2}\Omega\pi \quad \dots(1.3)$$

where

$$\Omega = \sum_{i=1}^m \delta_j + \sum_{j=1}^n E_j \gamma_j - \sum_{j=m+1}^q F_j \delta_j - \sum_{j=n+1}^p \gamma_j > 0 \quad \dots(1.4)$$

The series representation of the \bar{H} -function is given by [6, p.271, Eqn.(6)]

$$\begin{aligned} \bar{H}_{P,Q}^{M,N} \left[z \left| \begin{matrix} (a_j, \alpha_j; A_j)_{1,N}, (a_j, \alpha_j)_{N+1,P} \\ (b_j, \beta_j)_{1,M}, (b_j, \beta_j; B_j)_{M+1,Q} \end{matrix} \right. \right] \\ = \sum_{h=1}^M \sum_{r=0}^{\infty} \frac{\prod_{\substack{j=1 \\ j \neq h}}^M \Gamma(b_j - \beta_j s) \prod_{j=1}^N \{\Gamma(1 - a_j + \alpha_j s)\}^{A_j} (-1)^r z^s}{\prod_{j=M+1}^Q \{\Gamma(1 - b_j + \beta_j s)\}^{B_j} \prod_{j=N+1}^P \Gamma(a_j - \alpha_j s) r! \beta_h} \quad \dots(1.5) \end{aligned}$$

where

$$s = \xi_{h,r} = \frac{b_h + r}{\beta_h}$$

The conventional formulation may fail pertaining to the domain of quantum cosmology but Feynman path integrals [7,8]. Feynman path integrals reformulation of quantum mechanics is more fundamental than the conventional formulation in terms of operators. Feynman integrals are useful in the study and development of simple and multiple variable hypergeometric series which in turn are useful in statistical mechanics.

2. MAIN RESULTS

Here we shall evaluate the following integrals pertaining to M-series [9] with \bar{H} -function [7], a general class of polynomials [11].

(A)

$$\begin{aligned}
 & \int_0^1 \int_0^1 \left[\frac{(1-x)}{(1-xy)} y \right]^\tau \left[\frac{1-y}{1-xy} \right]^\psi \left[\frac{(1-xy)}{(1-x)(1-y)} \right] S_V^U \left[c_1 \left\{ \frac{(1-x)y}{(1-xy)} \right\}^{\lambda_1} \left\{ \frac{(1-y)}{(1-xy)} \right\}^{\theta_1} \right] \\
 & \cdot \bar{H}_{P,Q}^{M,N} \left[c_2 \left\{ \frac{(1-x)y}{(1-xy)} \right\}^{\lambda_2} \left\{ \frac{(1-y)}{(1-xy)} \right\}^{\theta_2} \right] {}_\alpha^M \left[c_3 \left\{ \frac{(1-x)y}{(1-xy)} \right\}^{\lambda_3} \left\{ \frac{(1-y)}{(1-xy)} \right\}^{\theta_3} \right] \\
 & \cdot \bar{H}_{p,q}^{m,n} \left[c_4 \left\{ \frac{(1-x)y}{(1-xy)} \right\}^{\lambda_4} \left\{ \frac{(1-y)}{(1-xy)} \right\}^{\theta_4} \right] dx dy \\
 & = \sum_{j=0}^{[V/U]} \sum_{h=1}^M \sum_{g,r=0}^{\infty} \frac{(-V)_{UJ}}{J!} A_{V,J} \frac{(-1)^r}{r!} \phi(s) c_1^j c_2^s c_3^g \frac{(u_1)_g \dots (u_\alpha)_g}{(v_1)_g \dots (v_\beta)_g} \frac{1}{\Gamma(\lambda g + 1)} \\
 & \cdot \bar{H}_{p+2,q+1}^{m,n+2} \left[c_4 \left| \begin{array}{l} (1-\tau-\lambda_1 J - \lambda_2 s - \lambda_3 g, \lambda_4; 1), (1-\psi-\theta_1 J - \theta_2 s - \theta_3 g, \theta_4; 1), (e_j, \gamma_j; E_j)_{1,n}, (e_j, \gamma_j)_{n+1,p} \\ (f_i, \delta_j)_{1,m}, (f_j, \delta_j; F_j)_{m+1,q}, \{1-\tau-(\lambda_1+\theta_1)J - (\lambda_2+\theta_2)s - (\lambda_3+\theta_3)g, (\lambda_4+\theta_4); 1\} \end{array} \right. \right] \dots (2.1)
 \end{aligned}$$

provided that

$$\operatorname{Re} \left(\tau + 1 + \lambda_2 \frac{b_j}{\beta_j} + \lambda_4 \frac{f_{j'}}{\delta_{j'}} \right) > 0, \quad \operatorname{Re} \left(\psi + 1 + \theta_2 \frac{b_j}{\beta_j} + \theta_4 \frac{f_{j'}}{\delta_{j'}} \right) > 0,$$

$$|\arg c_2| < \frac{1}{2}\Omega'\pi, \Omega' > 0, \quad |\arg c_4| < \frac{1}{2}\Omega\pi, \Omega > 0,$$

where $j=1, \dots, M; j'=1, \dots, m; (\tau, \psi, \lambda_1, \theta_1, \lambda_2, \theta_2, \lambda_3, \theta_3, \lambda_4, \theta_4) > 0$ and

$$\alpha \leq \beta, |c_3| < 1, c_1 > 0.$$

(B)

$$\begin{aligned} & \int_0^\infty \int_0^\infty f(x+y) x^{\tau-1} y^{\psi-1} S_V^U [c_1 x^{\lambda_1} y^{\theta_1}] \\ & \cdot \bar{H}_{P,Q}^{M,N} \left[c_2 x^{\lambda_2} y^{\theta_2} \left| \begin{array}{l} (a_j, \alpha_j; A_j)_{1,N}, (a_j, \alpha_j)_{N+1,P} \\ (b_j, \beta_j)_{1,M}, (b_j, \beta_j; B_j)_{M+1,Q} \end{array} \right. \right] {}_\alpha^M \beta [c_3 x^{\lambda_3} y^{\theta_3}] \\ & \cdot \bar{H}_{p,q}^{m,n} \left[c_4 x^{\lambda_4} y^{\theta_4} \left| \begin{array}{l} (e_j, \gamma_j; E_j)_{1,n}, (e_j, \gamma_j)_{n+1,p} \\ (f_j, \delta_j)_{1,m}, (f_j, \delta_j; F_j)_{m+1,q} \end{array} \right. \right] dx dy \\ & = \sum_{j=0}^{[V/U]} \sum_{h=1}^M \sum_{g,r=0}^\infty \frac{(-V)_{Uj}}{J!} A_{V,J} \frac{(-1)^r}{r!} \frac{\phi(s)}{\beta_h} \frac{(u_1)_g \dots (u_\alpha)_g}{(v_1)_g \dots (v_\beta)_g} \frac{1}{\Gamma(\lambda g + 1)} c_1^J c_2^s c_3^g \\ & \cdot \int_0^\infty f(z) z^{[\tau+\psi+(\lambda_1+\theta_1)J+(\lambda_2+\theta_2)s+(\lambda_3+\theta_3)g-1]} dz \\ & \times \bar{H}_{p+2,q+1}^{m,n+2} \left[c_4 z^{\lambda_4+\theta_4} \left| \begin{array}{l} (1-\tau-\lambda_1 J - \lambda_2 s - \lambda_3 g, \lambda_4; 1), (1-\psi-\theta_1 J - \theta_2 s - \theta_3 g, \theta_4; 1), (e_j, \gamma_j; E_j)_{1,n}, (e_j, \gamma_j)_{n+1,p} \\ (f_1, \delta_1)_{1,m}, (f_j, \delta_j; F_j)_{m+1,q}, \{1-\tau-\psi-(\lambda_1+\theta_1)J-(\lambda_2+\theta_2)s-(\lambda_3+\theta_3)g, (\lambda_4+\theta_4); 1\} \end{array} \right. \right] \end{aligned} \quad \dots(2.2)$$

provided that

$$\operatorname{Re} \left(\tau + \lambda_2 \frac{b_j}{\beta_j} + \lambda_4 \frac{f_{j'}}{\delta_{j'}} \right) > 0, \operatorname{Re} \left(\tau + \lambda_2 \left(\frac{a_j - 1}{\alpha_j} \right) + \lambda_4 \left(\frac{E_{j'}(e_{j'} - 1)}{\gamma_{j'}} \right) \right) < 0$$

$$\operatorname{Re} \left(\psi + \theta_2 \frac{b_j}{\beta_j} + \theta_4 \frac{f_{j'}}{\delta_{j'}} \right) > 0,$$

$$\operatorname{Re} \left(\psi + \theta_2 \left(\frac{a_j - 1}{\alpha_j} \right) + \theta_4 \left(\frac{E_{j'}(e_{j'} - 1)}{\gamma_{j'}} \right) \right) < 0$$

$$|\arg c_2| < \frac{1}{2}\Omega' \pi, \Omega' > 0,$$

$$|\arg c_4| < \frac{1}{2}\Omega \pi, \Omega > 0,$$

where $j = 1, \dots, M$; $j' = 1, \dots, m$; $(\tau, \psi, \lambda_1, \theta_1, \lambda_2, \theta_2, \lambda_3, \theta_3, \lambda_4, \theta_4) > 0$ and

$$\alpha \leq \beta, |c_2| < 1, c_1 > 0.$$

(C)

$$\begin{aligned} & \int_0^1 \int_0^1 f(xy)(1-x)^{\tau-1} (1-y)^{\psi-1} y^\tau S_V^U [c_1 (1-x)^{\lambda_1} (1-y)^{\theta_1} y^{\lambda_1}] \\ & \cdot \overline{H}_{P,Q}^{M,N} \left[c_2 (1-x)^{\lambda_2} (1-y)^{\theta_2} y^{\lambda_2} \left| \begin{array}{l} (a_j, \alpha_j; A_j)_{1,N}, (a_j, \alpha_j)_{N+1,P} \\ (b_j, \beta_j)_{1,M}, (b_j, \beta_j; B_j)_{M+1,Q} \end{array} \right. \right] {}_\alpha^\lambda M_\beta [c_3 (1-x)^{\lambda_3} (1-y)^{\theta_3} y^{\lambda_3}] \\ & \cdot \overline{H}_{P,Q}^{m,n} \left[c_4 (1-x)^{\lambda_4} (1-y)^{\theta_4} y^{\lambda_4} \left| \begin{array}{l} (e_j, \gamma_j; E_j)_{1,n}, (e_j, \gamma_j)_{n+1,p} \\ (f_j, \delta_j)_{1,m}, (f_j, \delta_j; F_j)_{m+1,q} \end{array} \right. \right] dx dy \\ & = \sum_{j=0}^{[V/U]} \sum_{h=1}^M \sum_{g,r=0}^{\infty} \frac{(-V)_{UJ}}{J!} A_{V,J} \frac{(-1)^r}{r!} \frac{\phi(s)}{\beta_h} \frac{(u_1)_g \dots (u_\alpha)_g}{(v_1)_g \dots (v_\beta)_g} \frac{1}{\Gamma(\lambda g + 1)} c_1^J c_2^s c_3^g \\ & \cdot \int_0^1 f(z) (1-z)^{\tau+\psi+(\lambda_1+\theta_1)J+(\lambda_2+\theta_2)s+(\lambda_3+\theta_3)g-1} dz \end{aligned}$$

$$\times \bar{H}_{p+2,q+1}^{m,n+2} \left[c_4 (1-z)^{(\lambda_4 + \theta_4)} \begin{matrix} (1-\tau - \lambda_1 J - \lambda_2 s - \lambda_3 g, \lambda_4; 1), (1-\psi - \theta_1 J - \theta_2 s - \theta_3 g, \theta_4; 1), (e_j, \gamma_j; E_j)_{1,n}, (e_j, \gamma_j)_{n+1,p} \\ (f_i, \delta_j)_{1,m}, (f_j, \delta_j; F_j)_{m+1,q}, \{1-\tau - \psi - (\lambda_1 + \theta_1)J - (\lambda_2 + \theta_2)s - (\lambda_3 + \theta_3)g, (\lambda_4 + \theta_4); 1\} \end{matrix} \right] \dots (2.3)$$

provided that

$$\operatorname{Re} \left(\tau + 1 + \lambda_2 \frac{b_j}{\beta_j} + \lambda_4 \frac{f_j}{\delta_j} \right) > 0, \operatorname{Re} \left(\lambda + \lambda_2 \frac{b_j}{\beta_j} + \lambda_4 \frac{f_j}{\delta_j} \right) > 0$$

$$\operatorname{Re} \left(\psi + \theta_2 \frac{b_j}{\beta_j} + \theta_4 \frac{f_j}{\delta_j} \right) > 0,$$

$$|\arg c_2| < \frac{1}{2}\Omega' \pi, \Omega' > 0,$$

$$|\arg c_4| < \frac{1}{2}\Omega \pi, \Omega > 0,$$

where $j = 1, \dots, M$; $j' = 1, \dots, m$; $(\tau, \psi, \lambda_1, \theta_1, \lambda_2, \theta_2, \lambda_3, \theta_3, \lambda_4, \theta_4) > 0$ and

$$\alpha \leq \beta, |c_3| < 1, c_1 > 0.$$

(D)

$$\int_0^1 \int_0^1 \left[\frac{(1-x)y}{1-xy} \right]^{\tau+\psi} \left[\frac{1-y}{1-xy} \right]^{\theta} \left[\frac{1}{(1-x)} \right] P_n(\tau, \theta) \left[1 - 2c_1 \left\{ \frac{y(1-x)}{1-xy} \right\}^{\lambda_1} \left\{ \frac{1-y}{1-xy} \right\}^{\theta_1} \right]$$

$$S_V^U \left[c_2 \left\{ \frac{y(1-x)}{(1-xy)} \right\}^{\lambda_2} \left\{ \frac{1-y}{1-xy} \right\}^{\theta_2} \right]$$

$$\bar{H}_{P,Q}^{M,N} \left[c_3 \left\{ \frac{y(1-x)}{1-xy} \right\}^{\lambda_3} \left\{ \frac{1-y}{1-xy} \right\}^{\theta_3} \begin{matrix} (a_j, \alpha_j; A_j)_{1,N}, (a_j, \alpha_j)_{N+1,P} \\ (b_j, \beta_j)_{1,M}, (b_j, \beta_j; B_j)_{M+1,Q} \end{matrix} \right]$$

$$\begin{aligned}
 & \cdot {}_{\alpha} M_{\beta}^{\lambda} \left[c_4 \left\{ \frac{y(1-x)}{1-xy} \right\}^{\lambda_4} \left\{ \frac{1-y}{1-xy} \right\}^{\theta_4} \right] \\
 & \cdot \bar{H}_{p,q}^{m,n} \left[c_5 \left\{ \frac{y(1-x)}{1-xy} \right\}^{\lambda_5} \left\{ \frac{1-y}{1-xy} \right\}^{\theta_5} \right|_{(f_j, \delta_j)_{1,m}, (f_j, \delta_j; F_j)_{m+1,q}}^{(e_j, \gamma_j; E_j)_{1,n}, (e_j, \gamma_j)_{n+1,p}} dx dy \\
 = & \sum_{j=0}^{[V/U]} \sum_{h=1}^M \sum_{g,r=0}^{\infty} \frac{(-V)_{UJ}}{J!} A_{V,J} \frac{(-1)^r}{r! \beta_h} \phi(s) \frac{(u_1)_g \dots (u_\alpha)_g}{(v_1)_g \dots (v_\beta)_g} \frac{1}{\Gamma(\lambda g + 1)} c_2^J c_3^s c_4^g \frac{(1+\tau)_n}{n!} \\
 & \cdot {}_2 F_1 \left[{}_{-n, \tau + \theta + n + 1}^{\quad 1 + \tau}; c_1 \right] \\
 & \cdot \bar{H}_{p+2,q+1}^{m,n+2} \left[c_5 \left| \begin{array}{l} (1-\tau-\psi-\lambda_1 k_1 - \lambda_2 J - \lambda_3 s - \lambda_4 g, \lambda_5; 1), (-\theta - \theta_1 k_1 - \theta_2 J - \theta_3 s, \theta_4 g, \theta_5; 1), (e_j, \gamma_j; E_j)_{1,n}, (e_j, \gamma_j)_{n+1,p} \\ (f_j, \delta_j)_{1,m}, (f_j, \delta_j; F_j)_{m+1,q}, \{-\tau - \theta - \psi - (\lambda_1 + \theta_1) k_1 - (\lambda_2 + \theta_2) J - (\lambda_3 + \theta_3) s - (\lambda_4 + \theta_4) g, (\lambda_5 + \theta_5); 1 \} \end{array} \right. \right]
 \end{aligned}$$

...(2.4)

provided that

$$\begin{aligned}
 & \operatorname{Re} \left(\tau + \psi + 1 + \lambda_3 \frac{b_j}{\beta_j} + \lambda_5 \frac{f_j}{\delta_j} \right) > 0, \quad \operatorname{Re} \left(\theta + 1 + \theta_3 \frac{b_j}{\beta_j} + \theta_5 \frac{f_j}{\delta_j} \right) > 0, \\
 & \cdot |\arg c_3| < \frac{1}{2} \Omega' \pi, \Omega' > 0, \quad |\arg c_5| < \frac{1}{2} \Omega \pi, \Omega > 0,
 \end{aligned}$$

where $j=1, \dots, M; j'=1, \dots, m; (\tau, \psi, \theta, \lambda_1, \theta_1, \lambda_2, \theta_2, \lambda_3, \theta_3, \lambda_4, \theta_4, \lambda_5, \theta_5) > 0$ and

$$\alpha \leq \beta, |c_4| < 1.$$

Proofs. The results (2.1) through (2.4) can be easily proved by using the same method as used by Chaurasia and Shekhawat [3].

3. PARTICULAR CASES

(3.A) By setting $A_j = B_j = 1, j=1,...,M; j=1,...,N$ in (2.1) through (2.4), we get the known results obtained by Chaurasia and Jagdev Singh ([2], eqn. (2.1), p.2, eqn.(2.2), p.2, eqn.(2.3), p.3, eqn. (2.4), p.4).

(3.B) By taking $\lambda = 1$, for the generalized hypergeometric function [9] and $A_j = B_j = 1, j = 1,...,M; j = 1,...,N$ in the results (2.1) through (2.4), we get the known results obtained by Chaurasia and Jagdev Singh ([2], eqn. (A.1) through (A.2), p.4,5).

(3.C) Using $A_j = B_j = 1, \lambda_1 \rightarrow 1, \theta_1 \rightarrow 0, \lambda_2 \rightarrow 0, \theta_2 \rightarrow 0, c_2 \rightarrow 1, \lambda_3 \rightarrow 0, \theta_3 \rightarrow 0, c_4 \rightarrow 1$ in equation (2.1), $A_j = B_j = 1, c_1 \rightarrow 0, \lambda_1 \rightarrow 1, \theta_1 \rightarrow 0, c_2 \rightarrow 1, \lambda_2 \rightarrow 0, \theta_2 \rightarrow 0, \lambda_3 \rightarrow 0$, in equation (2.2) and (2.3), we obtain the known results established by Chaurasia and Shekhawat ([3], eqn. 2.1` , p.184, eqn. (2.3), p.185, and eqn. (2.5), p.186).

ACKNOWLEDGEMENT

The authors are grateful to Professor H.M. Srivastava, University of Victoria, Canada for his kind help and valuable suggestions in the preparation of this paper.

REFERENCES

- [1] R.G. Buschman and H.M. Srivastava, “The \bar{H} -function associated with a certain class of Feymann integrals, J. Phys. A:Math. Gen. 23 (1990), 4707-4710.
- [2] V.B.L. Chaurasia and Jagdev Singh, “Certain integral properties pertaining to special functions”, SCIENTIA Series A: Mathematical Sciences, Vol.19 (2010), 1-6.
Copyright © 2012 SciResPub.

- [3] V.B.L. Chaurasia and A.S. Shekhawat, "Some integral properties of a general class of polynomials associated with Feynman integrals, Bull. Malays. Math. Sci. Soc. (2), 28(2) (2005), 183-189.
- [4] C. Fox, "The G and H-functions as symmetrical Fourier kernels, Trans. Amer. Math. Soc. 98 (1961), 395-429.
- [5] C. Grosche and F. Steiner, "Hand book of Feynman path integrals, Springer tracts in modern physics, Vol.145, Springer-Verlag, Berlin Heidelberg, New York (1993).
- [6] K.C. Gupta, Rashmi Jain and Aarti Sharma,"A study of unified finite integral transforms with application". J. Rajasthan Acad. Phy. Sci.2 (2003), 269-282.
- [7] A.A. Inayat-Hussain, "New properties of hypergeometric series derivable from Feynman integrals :I, Transformations and reduction formulae", J. Phys. A : Math. Gen. 20(1987), 4109-4117.
- [8] A.A. Inayat-Hussain, "New properties of hypergeometric derivable from Feynmann integrals : II. A generalization of the H-function", J.Phys. A: Math. (20) (1987), 4119-4128.
- [9] M. Sharma, "Fractional integration and fractional differentiation of the M-series", Frac. Cal.Appl. Anal., 11 (2008), 187-191.
- [10] P.Skibinski, "Some expansion theorems for the H-functions", Ann. Polon. Math., 23 (1970), 1-6.
- [11] H.M. Srivastava, "A contour integral involving Fox H-function", Indian J. Math. 14(1972), 1-6.
- [12] H.M. Srivastava and N.P. Singh, "The integration of certain products of the multivariable H-function with a general class of polynomials, Rend.Circ. Mat. Palermo 2(32) (1983), 157-187.
- [13] C. Szego,"Orthogonal polynomials, Amer. Math. Soc. Colloq. Publ. 23, Fourth edition, Amer. Math. Soc. Providence, Rhode Island (1975).