

Table (3): Clinical findings in patient groups.

Groups	Group(2) HCV (n=10)	Group(3) NASH (n=10)	Group(4) Cirrhosis (n=10)	Group(5) HCC (n=20)
Characteristics				
Jaundice (%)	0/10 (0%)	2/10 (20%)	4/10 (40%)	16/20 (80%)
Pallor (%)	1/10 (10%)	1/10 (10%)	3/10 (30%)	10/20 (50%)
GIT bleeding (%)	1/10 (10%)	1/10 (10%)	4/10 (40%)	8/20 (40%)
Blood transfusion (%)	1/10 (10%)	1/10 (10%)	2/10 (20%)	4/20 (20%)
Hepatic encephalopathy (%)	0/10 (0%)	0/10 (0%)	3/10 (30%)	10/20 (50%)
Ascites (%)	1/10 (10%)	0/10 (0%)	4/10 (40%)	10/20 (50%)
Liver size				
Average	10/10 (100%)	0/10 (0%)	0/10 (0%)	0/20 (0%)
Enlarged	0/10 (0%)	10/10 (100%)	3/10 (30%)	4/20 (20%)
Shrunken	0/10 (0%)	0/10 (0%)	7/10 (70%)	16/20 (80%)
Spleen size				
Average	7/10 (70%)	9/10 (90%)	3/10 (30%)	1/20 (5%)
Enlarged	3/10 (30%)	1/10 (10%)	5/10 (50%)	13/20 (65%)
Removed	0/10 (0%)	0/10 (0%)	2/10 (20%)	6/20 (30%)

