

Sensitivity and Specificity of PCR & Microscopy in detection of Babesiosis in domesticated cattle of Khyber Pakhtunkhwa, Pakistan.

Sumaira Shams¹, Sultan Ayaz^{1*}, Ijaz Ali², Sanuallah Khan¹, Irum Gul¹, Naila Gul¹ and Shahid Niaz Khan¹

¹Department of Zoology, Kohat University of Science and Technology Kohat Pakistan

²Institute of Biotechnology and Genetic Engineering, KP Agriculture University Peshawar Pakistan

Correspondence author:

Sultan Ayaz

Assistant Professor

Department of Zoology

Kohat University of Science and technology kohat Pakistan

Email; sultanayaz64@yahoo.com cell:+92-03348684196

ABSTRACT.

Background: Babesiosis is a tick-transmitted disease of veterinary and medical importance causes significant morbidity and mortality in cattle. This study is proposed to analyze/detect the active infection of Babesiosis in cattle by PCR and microscopic screening and to determine the sensitivity and specificity of the both diagnostic techniques in detection of Babesiosis in Khyber Pakhtunkhwa Pakistan **Methods:** Six hundred blood samples of clinically suspected cattle were collected from veterinary hospitals of district Karak and Kohat Khyber Pakhtunkhwa Pakistan. Thick and thin smear slides were examined under microscope. DNA was extracted from serum and was amplified by PCR using species specific primers. Amplified product was analyzed after electrophoresis under UV transilluminator. **Results:** Overall prevalence of *Babesia* was 27.5% (165/600) in cattle by PCR amongst these 20.6% (62/300) in calves and 34.3% (103/300) in cows. Similarly overall prevalence was 9.83% (59/600) by microscopy, amongst these 7.7% (23/300) in calves and 12% (36/300) in cows. Of these *B. bovis* was detected in 6.3% in calves, and 10.6% in cows and *B. bigemina* 4% in calves and 6.6% in cows by PCR while the mixed infection was 10.3% in calves and 17% in cows. The sensitivity of PCR was 100% and specificity was recorded 93.67% in detection of *B. bovis* as compare to Microscopy sensitivity was 36.84% and specificity 100% was observed. **Conclusions:** The study reveals that PCR is more sensitive technique in diagnosis against Babesiosis as compare to microscopy and recommended it for field application in Khyber Pakhtunkhwa Pakistan. So that proper preventive measure against this disease should be taken to increase the productivity of domesticated cattle.

Key Words: *Babesia*, cattle, PCR, Ticks, Khyber Pakhtunkhwa

1.INTRODUCTION

Babesiosis is economically the most important tick-borne disease of cattle (Bock et al., 2004), which causes extensive monetary loss to the cattle industry in the developing countries (Thammasirirak et al., 2003). The economic impact of babesiosis also occurs in other domestic animals, including horse, sheep, goats, pigs and dogs (Chaudry et al., 2010). Clinical signs of this disease are characterized by fever, anemia and hemoglobinuria in the infected cattle (Iseki et al., 2010) which may cause abortion in pregnant ones and reduced fertility in males especially bulls (Zulfiqar et al., 2012).

Babesiosis is caused by two intra-erythrocytic protozoan parasite of the genus *Babesia* (Family: *Babesiidae*)

namely *Babesia bovis* (*B. bovis*) and *Babesia bigemina* (*B. bigemina*) (Iseki et al., 2010). Generally *B. bovis* is more pathogenic than *B. bigemina* (Chaudry et al., 2010), however both can cause clinical disease and carrier infections (Goff et al., 2008). Ticks (*Boophilus* species) are the major vector for the transmission of both these spp. of *Babesia* (Chaudry et al., 2010), which are widely spread in tropical and subtropical countries particularly in Pakistan, Bangladesh and India as the environmental conditions favor the growth and development of many tick species (Irshad et al., 2010) *Babesia* in infected animals can be detected by microscopic examination of blood smears stained with Giemsa (Altay et al., 2008), but this method shows low sensitivity and time consuming (Adham et al., 2009). Serological methods (ELISA) may also be used but due to interspecies cross reactions, it is difficult to

differentiate between post exposure and present infection (Adham et al., 2009). Polymerase chain reaction (PCR), which is more sensitive and specific technique, offers an alternative approach for the diagnosis of babesiosis (Zulfiqar et al., 2012).

This study was aimed to determine the sensitivity and specificity of PCR and Microscopy diagnostic techniques in diagnosis of Babesiosis in the domesticated cattle in district Kohat and Karak of Khyber Pakhtunkhwa province Pakistan.

2. MATERIALS AND METHODS

2.1 sample collection.

Six hundred Blood samples were collected randomly from the clinically suspected cattle for babesiosis and were labeled the date of collection, sex, age and the areas where the sample were collected. The samples were transported in very sterilized container and stored at -20°C in Molecular Parasitology and Virology Laboratory Department of Zoology, Kohat University of Science and Technology Kohat Khyber Pakhtunkhwa Pakistan. Clinical data was recorded on designed proforma.

2.2 Microscopy

Thick and thin blood smears were fixed with absolute methanol for one minute and was stained by Giemsa stain. Stained slides were observed under 100x magnification using microscope (Olympus Japan) to detect *B. bigemina* and *B. bovis*. Pictures of positive slides were taken. *B. bigemina* is larger in size and having paired structure at an acute angle to each other. *B. bovis* is smaller in size and having paired form at an obtuse angle to each other. However it is hard to differentiate the two species by microscopic examination.

2.3 MOLECULAR DETECTION

2.4 DNA isolation

DNA was isolated from the whole blood using GF-1 Nucleic Acid isolation Kit (Vivantus USA) according to the manufacturer procedure.

2.5 DNA amplification and Detection

DNA was amplified for *B. bigemina* and *B. bovis* accordingly mentioned by Guido et al. (2002). Briefly, PCR program was started with initial denaturation at 94°C for 5 min, and 35 cycles of 94°C for 30 sec, 50°C for 30 sec, 72°C for 45 sec and final extension at 72°C for 10 minutes. Finally the amplified DNA was analyzed under UV transilluminator after electrophoresis on 1% agarose gel, and compared with 100bp DNA ladder (Fermentas USA), used as size marker.

2.6 Sensitivity and specificity of PCR and microscopy

Sensitivity values for PCR were obtained using the following correlations.

Sensitivity= true positive/ true positive + false negative X 100

Specificity as calculated as follows

Specificity = true negative/ true negative + false positive X 100 (Anderson et al., 1989).

3. RESULTS

The six hundred samples were randomly examined from the cattle population. The prevalence rate of Babesiosis was 27.5% (165/600) by PCR; among these 24% (72/300) prevalence was noted in district Karak and similarly 31% (93/300) in district Kohat (table.1)

The prevalence of Babesia species in cattle i.e calves and cows was recorded 20.6% (62/300) and 24.3% (103/300) through PCR technique respectively. Similarly 7.7% (23/300) and 12% (36/300) by microscopy technique was detected. In the present study the aetiological agent i.e. species of the Babesia were detected in cows and it was found that *B. bovis* 2.3% (7/300) followed *B. bigemina* 1.3% (4/300) and mixed infection 4% (12/300) through microscopy while *B. bovis* 6.3% (19/300), *B. bigemina* 4% (12/300) and mixed infection 10.3% (31/300) (table.2)

Similarly 300 random sampling were carried by both of the techniques in cows which shows *B. bovis* 3.68% (11/300), *B. bigemina* 2.3% (7/300) and mixed infection 6% (18/300) through microscopy while 10.6% (10/300), 6.6% (20/300) and 17% (51/300) were reported by PCR respectively (table3)

4. Discussion

Babesiosis is one of the most prevalent diseases, affecting cattle and causing huge losses to the cattle industry (Ziapour et al., 2011). The prevalence of babesiosis in cattle is about 1.2 billion in many countries of the world including Asia, Australia, Africa, South and Central America and the United States (Zulfiqar et al., 2012) while in Pakistan about 5.5- 42.8% cattle and buffaloes are exposed to this disease (Niazi et al., 2008). In order to decrease the monetary losses by the parasite it is mandatory to diagnose and treat Babesiosis well in time. The most commonly used methods are microscopic examination of blood smears stained with Giemsa and PCR which is more specific and sensitive technique for the detection of *Babesia* spp. present in blood of animals (Shahnawaz et al., 2011; Mesplet et al., 2011).

We found high prevalence, 27.5% of *Babesia* spp. (8.5% were positive for *B. bovis*, 5.33% was *B. bigemina* and mixed infection was 13.67% through PCR while 9.83% were positive by microscopic examination. Similarly results were found by other

researcher from different areas of Pakistan like Qadirabad (Chaudhry et al., 2010), Kasure (Durrani and Kamal 2008), North Iran (Ziapour et al., 2011), Brazil (Oliveira-Sequeira et al., 2005), Egypt (Adham et al., 2009), Tunisia (Ghirbi et al., 2008), Kalubya Governorate (Talkhan et al., 2010) and Southern Punjab (Zulfiqar et al., 2012). Moreover, small differences in the prevalence rate may be due to environmental factors of the areas, sample size and cattle breeds under study. Our results also showed that cows were more infected by *Babesia* spp as compared to calves which are in contrast with the findings of Niazi et al., (2008), Zulfiqar et al., (2012) Atif et al., (2012) who found more infections in calves in Punjab. This may be due to poor

managerial and environmental condition. More infestation of calves may be due to softer and thinner skin of young animals which help the vector (ticks) to transmit the disease easily. In the present study it was found that PCR was more sensitive than microscopy similar finding was reported in detection method of protozoa which depends on several factors including the species whether they are capable of human infection (Li et al., 2004) and adopted the microscopy procedure for the identification of protozoa by Ho et al., 1995; smith et al., 2002) while PCR identification was more sensitive than microscopic examination as reported by Jiang et al., 2005.

Table No.1 .Detection of Babesiosis in district Karak and Kohat Khyber Pakhtunkhwa, Pakistan by PCR

Districts	+ive Samples/total	Prevalence (%)
Kohat	93/300	31%
Karak	72/300	24%
Grand total	165/600	27.5%

Table 2: Prevalence of *B.bovis* and *B.bigemina* in Cattle by microscopy and PCR

Indicator	Calves(n=300)		Cows(n=300)	
	Blood smear (%)	PCR (%)	Blood smear (%)	PCR (%)
<i>B.bovis</i>	7(2.3%)	19(6.3%)	11(3.66%)	32(10.6%)
<i>B.bigemina</i>	4(1.3%)	12(4%)	7(2.3%)	20(6.6%)
<i>B.bovis/B.bigemina</i>	12(4%)	31(10.3%)	18(6%)	51(17%)
Grand total	23(7.7%)	62(20.6%)	12(4%)	103(34.3%)

Table No.3 sensitivity/ specificity of PCR and microscopic techniques against the detection of Babesiosis in domesticated cattle in Khyber Pakhtunkhwa Pakistan.

Parasites (n)	PCR			Microscopy		
	True+ive	Sensitivity%	Specificity%	True+ive	Sensitivity%	Specificity%
B.bovis	19	100	93.67	7	36.84	100
B.bigemina	12	100	96	4	25	100
B.bovis/B.bigemina	31	100	89.66	12	11.42	100

References cited.

Adham FK, Abd-el-Samie EM, Gabre RM, el-Hussein H (2009). Detection of tick blood parasites in Egypt using PCR assay I-Babesia bovis and Babesia bigemina. Parasitol Res., 105:721-730.

Altay K, Aydin MF, Dumanli N, Aktas M (2008). Molecular detection of Theileria and Babesia infections in cattle. Vet Parasitol., 158:295-301.

Atif FA, Khan MS, Iqbal HJ, Ali Z, Ullah S (2012). Prevalance of cattle tick infestation in three districts of the Punjab Pakistan. Pakistan J. of Sci., 64:49-53.

Bashir IN, Chaudhry ZI, Ahmed S,Saeed MA (2009). Epidemiological and vector identification studies on canine babesiosis. Pakistan Vet. J., 29(2): 51-54.

Bock R, Jackson L, de Vos A, Jorgensen W (2004). Babesiosis of cattle. Parasitol., 129: 247-69.

Chaudhry ZI, Suleman M, Younus M, Aslim A (2010). Molecular Detection of Babesia bigemina and Babesia bovis in Crossbred Carrier Cattle Through PCR. Pakistan J. Zool., 42(2): 201-204.

Durrani AZ, Kamal N (2008). Identification of ticks and detection of blood protozoa in friesian cattle by polymerase chain reaction test and estimation of blood parameters in district Kasur, Pakistan. Trop Anim Health Prod., 40: 441-447.

Ghirbi YM, Hurtado A, Brandika J, Khelif K, Ketata Z, Bouattour A (2008). A molecular survey of Theileria and Babesia parasites in cattle,with a note on the

distribution of ticks in Tunisia. Parasitol Res.,103: 435-442.

Ghosh S, Bansal GC, Gupta SC, Ray D, Khan MQ, Irshad H, Shahiduzzaman MD, Seitzer U, Jabbar S, and Ahmed (2007). Status of tick distribution in Bangladesh, India and Pakistan. Parasitol Res.,101: 207-216.

Goff, W.L, Johnson, W.C, Molloy, J.B, Jorgensen, W.K, Waldron, S.J, Figueroa, J.V, Matthee O, Adams DS, McGuire TC, Pino I, Mosqueda J, Palmer GH, Suarez CE, Knowles DP, McElwain TF (2008). Validation of a competitive enzyme-linked immunosorbent assay for detection of Babesia bigemina antibodies in cattle. Clin.Vaccine Immunol., 15: 1316–1321.

Guido FCL, Angela PS., Lloyd HL, Claudio RM (2002). Assessment of primers designed from the small ribosomal subunit RNA for specific discrimination between Babesia bigemina and B.bovis by PCR. Cien. Anim. Brasil., 3: 27-32.

Irshad N, Qayyum M, Hussain M, Qasim MK (2010). Prevalence of Tick Infestation and Theileriosis in Sheep and Goats. Pak Vet J., 30(3): 178-180.

Iseki H, Zhou Z, Kim C, Inpankaew T, Sununta C, Yokoyama N, Xuan X, Jittapalapong S, Igarashi I (2010). Seroprevalence of Babesia infections of dairy cows in northernThailand. Vet Parasitol., 170: 193-196.

Manan A, Khan Z, Ahmad B, Abdullah (2007). Prevalence and identification of Ixodid tick genera in frontier region Peshawar. J.of Agri and Bio Sci., 2:21-25.

Mesplet M, Palmer GH, Pedronic MJ, Echaide I, Christensen MF, Schnittger L, Lau AOT (2011). Genome-wide analysis of peptidase content and expression in a virulent and attenuated *Babesia bovis* strain pair. *Mol Biochem Parasitol.*, 179(2-2): 111–111
Muhammad G, Naureen A, Firyal S, Saqib M (2008). Tick control strategies in dairy production medicine. *Pakistan Vet. J.*, 28(1): 43-50.

Niazi N, Khan MS, Avais M, Khan JA, Pervez K, Ijaz M (2008). A study on babesiosis in calves at livestock experimental station Qadirabad and adjacent areas, Sahiwal, Pakistan. *Pak J Agri Sci.*, 45:13-16.

Oliveira-Sequeira TCG, Oliveira MC (Jr), Araujo JP, Amarante AF (2005). PCR-based detection of *Babesia bovis* and *Babesia bigemina* in their natural host *Boophilus microplus* and cattle. *Int J Parasitol.*, 35: 105–111.

Ramzan M, Khan MS, Avais M, Khan JA, Pervez K, Shahzad W (2008). Prevalence of ecto parasites and comparative efficacy of different drugs against Shahnawaz S, Ali M, Aslam MA, Fatima R, Chaudhry ZI, Hassan MU, Ali M, Iqbal F (2011). A study on the prevalence of a tick transmitted pathogen, *Theileria annulata*, and hematological profile in cattle from Southern Punjab (Pakistan). *Parasitol Res.*, 109(4): 1155-1160.

Soulby EJK (1982). *Helminths, Arthropods and Protozoa of domesticated animals*, 7th edition, Baillier Tindal. London UK.

Talkhan OFA, Radwan MEI, Ali MA (2010). Cattle babesiosis and associated biochemical alteration in Kalubya Governorate. *Nat Sci.*, 12: 24-27.

Thammasirirak S, Siriteptawee J, Sattayasai N, Indrakamhang P, Araki T (2003). Detection of *Babesia Bovis* in cattle by PCR-ELISA. *Southeast Asian J Trop Med Public Health.*, 34: 751-757.

Ziapour SP, Esfandiari B, Youssefi MR (2011). Study of the prevalence of babesiosis in domesticated animals with suspected signs in Mazandaran province, north of Iran, during 2008. *J Anim Vet Adv.*, 10: 712-714.

Zulfiqar S, Shahnawaz S, Ali M, Bhutta AM, Iqbal S, Hayat S, Qadir S, Latif M, Kiran N, Saeed A, Ali M, Iqbal F (2012). Detection of *Babesia bovis* in blood samples and its effect on the hematological and serum biochemical profile in large ruminants from Southern Punjab. *Asian Pacific Journal of Tropical Biomedicine.*, 104-108.
st tick infestation in cattle. *J. Anim. Pl. Sci.*, 18(1):17-19.

IJOART