

Stem cell therapy in oral and maxillofacial region:

RAYMOND BRITTO.P

Final year B.D.S

Dr.DANRAJ.M

Professor in Prosthodontics

SAVEETHA DENTAL COLLEGE

CHENNAI -77

ABSTRACT

Cells with unique capacity for self-renewal and potency are called stem cells. With appropriate biochemical signals stem cells can be transformed into desirable cells and the obtained literature on stem cell with respect to their properties, types and advantages of dental stem cells. Emphasis has been given to the possibilities of stem cell therapy in the oral and maxillofacial region including regeneration of tooth and craniofacial defects.[1]

Keywords: Oral and maxillofacial region, stem cells from exfoliated deciduous teeth, stem cell marker, tooth regeneration

INTRODUCTION

A classic stem cell should possess two properties namely self-renewal and potency.

Self-renewal is the capacity of the cell to undergo numerous cycles of cell division maintaining the undifferentiated state. An ideal stem cell should have the capacity of self renewal beyond the “Hayflicks” limit (the ability of

the cell to proliferate to about 40-60 population doublings before it achieves senescence).[2]

Potency means the differentiation capacity of the stem cells

STEM CELL TYPES

Stem cells can be broadly divided into

1. Embryonic stem cell
2. Adult stem cell
 - Hematopoietic stem cell
 - Mesenchymal stem cell
3. Induced pluripotent stem cell

Embryonic stem cell

Embryonic stem cells are capable of multipotential differentiation but clinical feasibility is limited due to ethical issues. The inner cell mass (the part that would form fetus) of the embryo is used to form embryonic cell lines. Embryonic stem cells has a potential to differentiate into germ layers namely ectoderm, endoderm and mesoderm. Tumorigenesis and immune rejection is common with embryonic stem cells.[3]

Adult stem cell

Adult stem cells are multipotent stem cells. They have been harvested from different kind of tissues like bone marrow, umbilical cord, amniotic fluid, brain tissue, liver, pancreas, cornea, dental pulp, and adipose tissue. Adult stem cells are comparatively easier to isolate and do not have any ethical issues. Immune rejection and teratoma formation is also rare with adult stem cells. Adult stem cells are commonly used in current day practice.[4]

Induced pluripotent stem cell

Induced pluripotent stem cells(IPS) is an evolving concept in which 3–4 genes found in the stem cells are transfected into the donor cells using appropriate vectors. The stem cells thus derived by culturing will have

properties almost like embryonic stem cells. This path breaking discovery may have a major role in future stem cell therapy.[5]

SOURCES OF STEM CELLS

The oral and maxillofacial region can be treated with stem cells from the following sources

1. Adipose tissue
2. Stem cells from oral and maxillofacial region

Bone marrow

Bone marrow stem cells (BMSCs) can be harvested from sternum or iliac crest. It is composed of both hematopoietic stem cells and mesenchymal stem cells (MSCs). The majority of oro-maxillofacial oral structures are formed from mesenchymal cells. The advantage of bone marrow is that it has a larger volume of stem cells and can be differentiated in to wide variety of cells. Isolation of BMSCs can be carried out only under general anesthesia with possible post operative pain.[6]

Adipose tissue

They can be harvested from the lipectomy or liposuction aspirate. Adipose derived stem cells (ADSCs) contain a group of pluripotent mesenchymal stem cells that exhibit multilineage differentiation. Advantage of adipose tissue is that it is easily accessible and abundant in many individuals.[7]

Stem cells from the oro-maxillofacial region

Stem cells from oral and maxillofacial region predominantly contain mesenchymal stem cells. In oral and maxillofacial area different types of dental stem cells were isolated and characterized. They include

- Dental pulp stem cells (DPSCs)
- Stem cells from exfoliated deciduous teeth (SHED)
- Periodontal ligament stem cells (PDLSCs)
- Stem cells from apical papilla (SCAP)

- Dental follicle progenitor cells (DFPCs)

These dental stem cells have MSC like qualities, such as self-renewal and differentiation potential. [8]

DENTAL STEM CELL ADVANTAGES

The advantages of stem cells from oral and maxillofacial region is that

1. Have high plasticity.
2. It can be cryopreserved for longer period (Ideal for stem cell banking).
3. It showed good interaction with scaffold and growth factors.
4. Stem cells transplantations can cause pathogen transmission and also need immunosuppression, so autologous stem cell source is the best option. Dental pulp stem cells will be better fitting tool due to easy surgical access, the very low morbidity of the anatomical site after the collection of the pulp[9].

STEM CELL THERAPY IN THE ORO-MAXILLOFACIAL REGION

The structures of interest in oral and maxillofacial region include the enamel, dentin, dental pulp, cementum, periodontal ligament, craniofacial bones, the temporo mandibular joint, ligaments, skeletal muscles, tendons, skin, subcutaneous soft tissue, and salivary glands.

Regeneration of dentin, pulp

Dental pulp tissue has the regenerative potential to form dentin in response to any injury. Tubular dentin formation was observed when human pulp stem cells with scaffold (hydroxyapatite/tricalcium phosphate)

Regeneration of the pulp inside the damaged tooth can be the basic clinical application of stem therapy in dentistry. Root canal treatment in a young permanent molar will stop the tooth's continuous maturation process there by leaving thin egg shell like weak tooth that is susceptible to fracture.

Regeneration of pulp with stem cell therapy will be a better option. Stem cells harvested from the pulp of unwanted teeth like third molar can be

utilized to regenerate the pulp of severely injured tooth there by preventing the need for endodontic treatment in adults.[10]

STEM CELLS IN PERIODONTAL REGENERATION

Stem cells will be a promising tool for regenerating the periodontal structures such as periodontal ligament and other supporting elements.. BMSCs have the ability to produce alveolar bone, periodontal ligament,. Thus, BMSCs provides an alternative source for the treatment of periodontal diseases. Autologous mesenchymal stem cells from iliac crest in combination with platelet rich plasma from peripheral blood was used for periodontal regeneration. Significant closure of bone defect and improvement of attachment level was observed after one year . It also showed good healing and regeneration of interdental papilla.

Iwata *et al.* harvested and expanded primary canine PDL cells *in vitro* and also made into transplantable constructs containing PGA Scaffold and PDL cell sheets. The transplantable constructs in combination with porous bTCP(b –tricalcium phosphate) induced regeneration of periodontal structures, including alveolar bone, cementum, and periodontal fibers.[11]

Regeneration of craniofacial defects

Stem cells can be useful in the regeneration of bone and to correct large craniofacial defects due to cyst enucleation, tumor resection, and trauma. The closure of a bone defect is commonly carried out with the transfer of tissue, which have disadvantages like, not able to restore the unique function of the lost part, donor site morbidity, accompanied by scarring, infection and loss of function. Adipose derived stem cells was used to treat the calvarial defect (120 cm²) of a 7-year-old girl who had severe head injury.

Autologous adipose stem cells were extracted from gluteal region along with iliac crest bone graft. Autologous fibrin glue that holds the cells in place was prepared by cryoprecipitation. This successful technique has given new rays of hope that ADSCs can be used for difficult reconstructive procedures.

Soft tissue reconstruction in the oromaxillofacial region is of paramount importance when there is significant loss of soft tissues during surgery or trauma. Various methods including graft and flap transfer has been tried that produced donor site morbidity. Adipose cells with appropriate shaped scaffold can be used for reconstruction of soft tissues.

Stem cells isolated from dental pulp has a potential to differentiate into osteoblasts and are a good source for bone formation. Stem cells from oral and maxillofacial region can be combined with bone marrow stem cells to correct larger defects. up bone defects. This new procedure has added advantage of permitting the transplantation of more cells and better integrity compared with cell suspensions or gels.[12]

Future tissues

Future tissues like tissue engineered bone grafts, engineered joints and cranial sutures can be developed with stem cell therapy. A team of professionals including stem cell biologists, molecular biologists, geneticists, polymer and materials scientists, mechanical engineers and clinicians with knowledge of oral and maxillofacial disorders is needed to develop the field of craniofacial tissue engineering. The ability to design anatomically viable and functional bone would have great potential for oromaxillofacial reconstructions of congenital defects, cancer resections, and trauma. The anatomically shaped viable bone grafts like articular condyles can be engineered by using adult mesenchymal stem cells and biomimetic scaffold bioreactor.

Tissue engineered temporo mandibular joint was created by having natural bone building process as an inspiration. Condyle shaped scaffolds were made using decellularized bone with help of digitized clinical images. Stem cells were seeded into the scaffold and placed in a bioreactor chamber containing culture medium. In future this technique can be applied to regenerate other bones in oromaxillofacial region.[13]

Tooth regeneration

The regeneration of adult teeth will be possible in future with the newer advancement in stem cell therapy and tissue engineering. Regenerative procedures would be better fitting and alternative tool in place of dental implants. The tooth crown structure can be regenerated using tissue engineering techniques that combine stem cells and biodegradable scaffolds. Epithelial mesenchymal interactions are mandatory in tooth development. “These interactions are characterized by the reciprocal exchange of signals between these two naïve germ layer tissues and result in the emergence of unique terminal phenotypes with their supporting cells”.[14]

Tooth regeneration involves three key elements which include

- Inductive morphogenes
- Stem cells
- Scaffold

Steps involved in regeneration of tooth are

1. Harvesting and expansion of adult stem cells.
2. Seeding the stem cells into scaffold which provides optimized environment.
3. Cells are instructed with targeted soluble molecular signals spatially.
4. Confirming the gene expression profile of the cells for next stage in odontogenesis.[15]

CONCLUSION

The future dentistry will be more of regenerative based, where patients own cells can be used to treat diseases. Stem cell therapy has got a paramount role as a future treatment modality in dentistry. Regenerative dentistry will have to go in pace with regenerative medicine. On the other hand, stem cells should be differentiated to the appropriate cell types before they can be used clinically, otherwise it might lead to deleterious effects. Determining the role of local conditions such as the type of scaffold and the presence of the microorganisms should be very carefully analyzed. Longer patient follow up is needed to study the life time of regenerated tissue.

REFERENCES

1. Fortier LA. Stem cells, classifications, controversies and clinical applications. *Vet Surg.* 2005;34:415–23. [[PubMed](#)]
2. Bluteau G, Luder HU, De Bari C, Mitsiadis TA. Stem cells for tooth engineering. *Eur Cell Mater.* 2008;16:1–9. [[PubMed](#)]
3. Gardner RL. Stem cells: Potency, plasticity and public perception. *J Anat.* 2002;200:277–82. [[PMC free article](#)] [[PubMed](#)]
4. Hayflick L. The limited *in vitro* lifetime of human diploid cell strains. *Exp Cell Res.* 1965;37:614–36. [[PubMed](#)]

5. Gurdon JB, Byrne JA. The first half-century of nuclear transplantation. *Proc Natl Acad Sci U S A*. 2003;100:8048–52. [[PMC free article](#)] [[PubMed](#)]
6. Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS, et al. Embryonic stem cell lines derived from human blastocysts. *Science*. 1998;282:1145–7. [[PubMed](#)]
7. Keller GM. *In vitro* differentiation of embryonic stem cells. *Curr Opin Cell Biol*. 1995;7:862–9. [[PubMed](#)]
8. Wu DC, Boyd AS, Wood KJ. Embryonic stem cell transplantation: Potential applicability in cell replacement therapy and regenerative medicine. *Front Biosci*. 2007;12:4525–35. [[PubMed](#)]
9. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*. 2006;126:663–76. [[PubMed](#)]
10. Zuk PA, Zhu M, Mizuno H, Huang J, Futrell JW, Katz AJ, et al. Multilineage cells from human adipose tissue: Implications for cell- based therapies. *Tissue Eng*. 2001;7:211–28. [[PubMed](#)]
11. Gronthos S, Mankani M, Brahimi J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) *in vitro* and *in vivo*. *Proc Natl Acad Sci U S A*. 2000;97:13625–30. [[PMC free article](#)] [[PubMed](#)]
12. Miura M, Gronthos S, Zhao M, Lu B, Fisher LW, Robey PG, et al. SHED: Stem cells from human exfoliated deciduous teeth. *Proc Natl Acad Sci U S A*. 2003;100:5807–12. [[PMC free article](#)] [[PubMed](#)]
13. Seo BM, Miura M, Gronthos S, Bartold PM, Batouli S, Brahimi J, et al. Investigation of multipotent postnatal stem cells from human periodontal ligament. *Lancet*. 2004;364:149–55. [[PubMed](#)]
14. Sonoyama W, Liu Y, Yamaza T, Tuan RS, Wang S, Shi S, et al. Characterization of the apical papilla and its residing stem cells from human immature permanent teeth: A pilot study. *J Endod*. 2008;34:166–71. [[PMC free article](#)] [[PubMed](#)]

15. Morsczeck C, Götz W, Schierholz J, Zeilhofer F, Kühn U, Möhl C, et al. Isolation of precursor cells (PCs) from human dental follicle of wisdom teeth. *Matrix Biol.* 2005;24:155–65. [[PubMed](#)]

IJOART