

TO ASSESS THE EFFECT OF MATERNAL BMI ON OBSTETRICAL OUTCOME

Dr.Shuchi LAKHANPAL, M.B.B.S, Dr Asha AGGARWAL, M.D., Dr. Gurcharan KAUR , M.D.

Place of Study – Department of Obstetrics and Gynecology, Kasturba Hospital, Delhi, India

ABSTRACT:

AIMS: To assess the effect of maternal BMI on complications in pregnancy, mode of delivery, complications of labour and delivery.

METHODS:

A crosssectional study was carried out in the Obst and Gynae department, Kasturba Hospital, Delhi. The study enrolled 100 pregnant women. They were divided into 2 groups based on their BMI, more than or equal to 30.0 kg/m² were categorized as obese and less than 30 kg/m² as non obese respectively. Maternal complications in both types of patients were studied.

RESULTS:

CONCLUSION: As the obstetrical outcome is significantly altered due to obesity, we can improve maternal outcome by overcoming obesity. As obesity is a modifiable risk factor, preconception counseling creating awareness regarding health risk associated with obesity should be encouraged and obstetrical complications reduced.

KEY WORDS: BMI, obesity, obstetrical outcome, preeclampsia, casaerean section

INTRODUCTION

WHO describes obesity as “One of the most blatantly visible, yet most neglected, public health problems that threaten to overwhelm both more and less developed countries”. Obesity is a major public health issue and as per WHO, it is a “killer disease” at par with HIV and malnutrition. Even in countries like India, significant proportion of overweight and obese coexist

with the undernourished. Lifestyle modifications over the years have led to a more sedentary lifestyle. This is of global concern,¹ as excess bodyweight is now the sixth important risk factor contributing to disease worldwide and increased level of obesity may result in a decline in life expectancy in the future.²

The body mass index (BMI), or Quetelet index, is a heuristic proxy for human body fat based on an individual's weight and height. It was devised between 1830 and 1850 by the Belgian polymath Adolphe Quetelet during the course of developing "social physics".³

Obesity in pregnant women is associated with increased risk of Gestational diabetes, thromboembolism and is associated with hyperlipidemia and preeclampsia.

Obese women are more likely to undergo induction of labour, failed induction, operative vaginal delivery, shoulder dystocia and third and fourth degree perineal lacerations. Frequency of both 'Elective' and 'Emergency' caesarean section is increased in obese women. Anaesthetic complications like failed regional blocks and difficult intubation are more common in obese women. Also, there is an increased number of large for gestational age infants, lower apgar score and gross congenital malformations.

RESEARCH ELABORATIONS

MATERIALS N METHODS

Place of study - Deptt. Of Obstetrics and Gynaecology, Kasturba hospital, Delhi

Sample size - 100. 50 in each of the 2 groups (divided on the basis of BMI)

Study period - 1 April 2011- 20 April 2012

Type of study - Comparative Prospective study.

Statistical method used -

The data collected during the study is presented in the tabular form along with appropriate graphs and charts to draw meaningful observations and interpretations. Wherever deemed necessary, suitable statistical techniques are applied to establish the cause and effect relationships between selected variables. The differences in statistical parameters for different

outcomes of pregnant women with BMI>30 were tested statistically using appropriate tests viz. t-test, Fisher exact test, Chi square tests etc and the results are presented with p values < 5% considered statistically significant.

BMI formula

The BMI is equal to a person's weight divided by their height .It is calculated either as;

$$\text{BMI} = (\text{weight in pounds/ height in inches}) \times 703$$

Or

$$\text{BMI} = (\text{weight in kilograms /height in meters}^2)$$

Based on this, patients to be studied will be divided into 2 groups of 50 patients each –

1. BMI less than 30
2. BMI more than 30

Inclusion criterion –

1. Primigravida with singleton pregnancy
2. Patients with gestational age more than 28 weeks

Exclusive criterion

1. Multifetal gestation
2. Multigravida

MATERIALS

The weighing machine used was from Equinox, an electronic personal scale CE.

Model : EB 1003

Strain gauge sensor

Capacity : 150kg(33016/24 stone)

Division : 0.1kg(0.216)

1.0”(25 mm) LCD digits

Low battery/ overload indication

Power : 1pc*3 V lithium cells (CR 2032)

Stadiometer used was from Bio Plus. A height measuring tape

Model no : 26M/1013522

Model approval mark : IND/09/2005/815

Size : 200cm / 78 inch

METHODOLOGY

Pregnant women coming for admission to labour room at the time of delivery were enrolled in the study after informed consent. A complete history work up and examination was done for the patient.

HISTORY

In all cases detailed history of the patient was taken including

.Name, age, education, religion, socio economic status

.Presenting complaints – Labour pains. Leaking per vaginum. hypertension, DM,

.History of present illness – if any

.Menstrual History – Last menstrual period, age of menarche, duration of Cycle, Length of cycle, Blood loss

.Obstetric History – Gravida, Parity, Number of live issues

.Past History, medical and surgical History – Any associated medical condition like diabetes mellitus, hypertension, tuberculosis, thyroid disease, asthma, any previous surgery.

.Family history- especially for obesity, diabetes and hypertension.

EXAMINATION

General examination- including general condition, hydration, PR, BP, temperature, pallor, icterus, cyanosis, edema, JVP, LN.

Weight(in kgs) was measured in kilograms. Patients were weighed without shoes, wearing light indoor clothes.

Height(in metres) was measured using a stadiometer. The patients were made to stand erect on the floor barefoot with both ankles together and parallel to each other. The head of the patient was held in such a position that the line joining the tragus and outer canthus of eye were in a horizontal plane (Frankfurts Plane), with the individual standing straight next to the wall with the heels, buttocks, shoulders and occiput touching the wall. The data were used to calculate Quetelet index or the BMI using the formula $BMI = \text{weight (kg)} / \text{height}^2 (\text{in m})$.

Systemic examination including cardiovascular, respiratory, central nervous system to rule out any systemic pathology

Per abdomen examination including contour, distension, venous prominence, stria, fundal height, presentation, fetal heart rate, regularity, estimated liquor, fetal weight, head floating/engaged. Also, local examination including vulva, vagina, urethra and Per speculum examination for cervix and vagina. Detailed Per vaginal examination was done for dilatation, effacement, position of cervix, station of presenting part, BISHOPS Scoring of the patient was then done. We also saw for adequacy of pelvis, leaking per vaginum/bleeding per vaginum.

INVESTIGATIONS

1. Blood group
2. CBC, ESR
3. FBS, PPBS
4. VDRL, HIV
5. Urine routine and microscopy
6. Obstetrical ultrasonography
7. Any other investigation needed as per patients requirement

After detailed history and examination, and after fulfilling the criterion for inclusion in the study, patients were divided into 2 groups-

1. BMI less than 30
2. BMI more than 30

In both the groups, fetomaternal outcome was studied along the following lines-

1. PREGNANCY ASSOCIATED CONDITIONS like hypertension, diabetes mellitus, abnormal presentations, IUGR, prematurity, postmaturity, any other illness
2. MODE OF DELIVERY – Normal vaginal delivery elective or emergency casaerean section, instrumental delivery.
3. LABOUR AND DELIVERY OUTCOME- Spontaneous or induced labour. First stage was studied to see progress of labour, and any complication like fetal distress, incoordinate uterine contractions, non progress of labour. Second stage to be studied for mode of delivery and any other complication, third stage for tear/PPH or any other complication.

4. CASAEREAN OUTCOME- difficulty in opening abdomen, uterine atony and any other complication.

ETHICAL ISSUES

As this was an observational study with no unethical interventions, or danger to the patient due to the study itself, it is an ethically sound study. Ethical clearance was taken by the hospital committee for the same.

RESULTS

A total of 100 cases, 50 with BMI>30 and 50 with BMI<30 were included in this study undertaken at Kasturba Hospital, Delhi. The primigravidas who presented in the labour room after 28 weeks of gestation were included. The antenatal, intrapartum, postpartum and neonatal assessment was done and outcome of each pregnancy in terms of maternal and perinatal morbidity and mortality were studied.

1.AGE DISTRIBUTION AND ITS RELATION WITH BMI

6% patients in the BMI >30 category were less than 20 years of age, 46% were in the 21-25 years age category, 34% in 26-30 and 14% in the 31-35 years of age. Also, in the BMI <30 category, 16% women were less than 20 yrs of age, 56% in 21-25 years, 24% in 26-30 years of age and only 4% in the 31-35 years. Mean age was 25.92 in the BMI>30 group compared with 24.2 in the BMI<30 group. We conclude that 48% of the BMI >30 category women were >26 years of age, whereas only 28% of the BMI < 30 group were in the >26 years category. If we consider patients above and below 25 years of age in different BMI categories, the p value comes out to be 0.039 making the difference statistically significant.

TABLE NO. 1 AGE DISTRIBUTION AND ITS RELATION WITH BMI

AGE	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE
<20	8	16	3	6	P=0.039 K ² =4.24
21-25	28	56	23	46	
26-30	12	24	17	34	
31-35	2	4	7	14	

2.ANTEPARTUM COMPLICATIONS AND ITS RELATION WITH BMI

Out of a total of 50 pregnancies in each category, only 32% patients in the BMI>30 category were free of complications and the number increased to 78% when the BMI was less than 30.

Preeclampsia complicated 8% of the pregnancies with BMI <30 and 38% of the patients with BMI>30 obese. The difference was statistically significant with a p value of 0.0003.

Eclampsia was found in 2% patients in the BMI >30 category, and was not found in BMI <30 category. P value of 1 was statistically insignificant.

Retinopathy was 6% in the BMI >30 category and 2% in BMI <30. The difference was statistically insignificant with a p value of 0.617.

Also, GDM complicated 2% of the pregnancies with BMI < 30 and 6% in the BMI >30 category. The difference was statistically insignificant with a p value of 0.617.

IUGR was present in 4% of the pregnancies with BMI < 30 and 6% in the BMI >30 category. The difference was statistically insignificant with a p value of 0.646.

Preterm labour pains occurred in 6% of the pregnancies with BMI < 30 and 10% in the BMI >30 category. The difference was statistically insignificant with a p value of 0.54.

TABLE NO. 2 ANTEPARTUM COMPLICATIONS AND ITS RELATION WITH BMI

COMPLICATION	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE D.F=1
PRECLAMPSIA	4	8	19	38	P=0.0003 K ² =12.7
ECLAMPSIA	0	0	1	2	F.P=1 K ² =1.01
RETINOPATHY	1	2	3	6	F.P=0.617 K ² =1.04
GDM	1	2	3	6	F.P=0.617 K ² =1.04
IUGR	2	4	3	6	P=0.646 K ² =0.21,
PRETERM	3	6	5	10	K ² =0.54 F.P=0.7149
NO COMPLICATION	39	78	16	32	
TOTAL	50	100	50	100	

3. MALPRESENTATIONS AND ITS RELATION WITH BMI

Malpresentations were present in 2% patients with BMI <30 category and 4% in the BMI>30.

The difference was statistically insignificant with a p value of 1.

TABLE NO. 3 MALPRESENTATIONS AND ITS RELATION WITH BMI

MALPRESENTATION	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE D.F=1
NORMAL	49	98	48	96	F.P=1 K ² =0.34
ABNORMAL	1	2	2	4	

4. PERIOD OF GESTATION (POG) AT DELIVERY AND ITS RELATION WITH BMI

Preterm labour pains were present in 6% of the BMI < 30 group and 10% in BMI > 30 category.

The difference was statistically insignificant with a p value of 0.7149.

Mothers reaching beyond term (post term) were 4% in the BMI < 30 group and no posterm patients were seen in the BMI > 30 group. The difference was statistically insignificant with a p value of 0.4949.

TABLE NO. 4 PERIOD OF GESTATION (POG) AT DELIVERY AND ITS RELATION WITH BMI

POG	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE D.F=1
PRETERM (<37 weeks)	3	6	5	10	F.P=0.7149 K ² =0.54
TERM	45	90	45	90	
POST-TERM	2	4	0	0	F.P=0.4949

(>40 weeks)					$K^2=2.04$
-------------	--	--	--	--	------------

5. INTRAPARTUM COMPLICATIONS AND ITS RELATION WITH BMI

Fetal distress was present in 6% patients with BMI <30 category and was absent in the BMI>30 group. The difference was statistically insignificant with a p value of 0.24.

Also, NPOL was present in 2% patients with BMI <30 category and was absent in the BMI >30. The difference was statistically insignificant with a p value of 1.

Also, failure of induction occurred in 2% patients with BMI <30 and in 2% with BMI>30. No statistical analysis could be done due to similar values and it was found at equal frequency in both the groups.

Shoulder dystocia was present in only 2% of the patients in the BMI>30 category, whereas it was absent in patients with BMI<30. The difference was statistically insignificant with a p value of 1.

TABLE NO. 5 INTRAPARTUM COMPLICATIONS AND ITS RELATION WITH BMI

COMPLICATION	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE D.F=1
FETAL DISTRESS	3	6	0	0	K ² =3.09, F.P=0.24
NPOL	1	2	0	0	K ² =1.01, F.P=1
FAILED INDUCTION	1	2	1	2	No statistical analysis
SHOULDER DYSTOCIA	0	0	1	2	K ² =1.01, F.P=1

NONE	44	90	48	96	
------	----	----	----	----	--

6. MODE OF DELIVERY AND ITS RELATION WITH BMI

Mode of delivery was normal vaginal delivery in 76% of the BMI <30 category patients and 44% in BMI >30 category. One patient (2% patients) in the BMI<30 group required forceps for delivery of baby. Casaerean sections were required in 22% patients in BMI<30 category and in 56% patients in BMI>30 category. The difference was statistically significant with a p value of <0.001.

TABLE NO. 6 MODE OF DELIVERY AND ITS RELATION WITH BMI

MODE OF DELIVERY	BMI<30	BMI>30	K ² AND P VALUE
NORMAL	38(76%)	22(44%)	K ² = 12.15, D.F=1, p-value < 0.001
INSTRUMENTAL	1(2%)	0(0%)	
LSCS	11(22%)	28(56%)	

7. ANAESTHETIC COMPLICATIONS AND ITS RELATION WITH BMI

Anaesthetic complications including failed attempt at spinal anaesthesia and resort to general anaesthesia and intraoperative ECG changes of T wave inversion an ST segment depression were seen in the patients. These occurred in none of the patients in BMI <30 category and in 10.17% patients undergoing LSCS in BMI>30 category. Statistical analysis revealed that p value was 0.545 making the difference insignificant.

TABLE NO. 7 ANAESTHETIC COMPLICATIONS AND ITS RELATION WITH BMI

ANAESTHETIC	BMI<30(%)	BMI>30(%)	K ² AND P
-------------	-----------	-----------	----------------------

COMPLICATIONS			VALUE
FAILED SPINAL	0(0%)	2(6.78%)	K ² = 1.28, D.F=1, F.P= 0.545
ECG CHANGES	0(0%)	1(3.39%)	
NONE	11(100%)	25(89.83%)	
TOTAL	11(100%)	28(100%)	

8. NEED FOR LSCS AND ITS RELATION WITH BMI

In the BMI<30 group, 27.27% patients had an elective LSCS whereas 72.72% had an emergency LSCS. In BMI>30 group, 35.714% patients had an elective LSCS whereas 64.285% had an emergency LSCS. The results were statistically insignificant with a p value of 0.719.

TABLE NO. 8 NEED FOR LSCS AND ITS RELATION WITH BMI

LSCS	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE
ELECTIVE	3	27.27	10	35.714	K ² =0.25 F.P=0.719 D.F.=1
EMERGENCY	8	72.72	18	64.285	

TOTAL	11	100	28	100

9. LSCS SURGICAL COMPLICATIONS AND ITS RELATION WITH BMI

Intraoperatively, we found that lower segment caesarean sections in BMI>30 group had higher incidence of bladder injury/ difficulty in opening/ trauma to neighbouring structures. In 2% patients with BMI <30 category and 10% in BMI>30 category, intraoperative LSCS complications were seen. Statistical analysis showed that $K^2 = 2.84$ and p value = 0.204 making the difference statistically insignificant.

The complications included bladder injury in the BMI<30 patient (2%). Broad ligament rent was seen in 1 patient in BMI>30. We experienced difficulty in opening the abdomen for LSCS in 4 patients in the BMI>30 group, making a total 10% complication rate in the BMI>30 group.

TABLE NO. 9 LSCS SURGICAL COMPLICATIONS AND ITS RELATION WITH BMI

SURGICAL COMPLICATIONS OF LSCS	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE D.F=1
INTRAOPLSCS	1	2	5	10	K ² =2.84, P=0.204
NO COMPLICATION	49	98	45	90	
TOTAL	50	100	50	100	

10. MODE OF TERMINATION OF PREGNANCY AND ITS RELATION WITH BMI

Inductions were done in 12% of the BMI <30 category and 14% of the the BMI >30 category.

The difference was statistically insignificant with a p value of 0.766.

TABLE NO. 10 MODE OF TERMINATION OF PREGNANCY AND ITS RELATION WITH BMI

LABOUR	BMI < 30	BMI > 30	K ² AND P VALUE
NO. OF PATIENTS INDUCED	6(12%)	7(14%)	K ² =0.09 P=0.766
SPONTANEOUS LABOUR	41(82%)	33(66%)	
ELECTIVE LSCS	3(6%)	10(20%)	

11. POSTPARTUM COMPLICATIONS (VAGINAL DELIVERY) AND ITS RELATION WITH BMI

PPH occurred in 2% of the patients with BMI <30 category and in 4% of the patients in the BMI>30 group. The difference was statistically insignificant with a p value of 0.604.

Cervical/ Paravaginal tears were present in 2% of the BMI <30 category and 4% in BMI>30 category. The difference was statistically insignificant with a p value of 0.604.

TABLE NO. 11 POSTPARTUM COMPLICATIONS (VAGINAL DELIVERY) AND ITS RELATION WITH BMI

COMPLICATION	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE D.F=1
PPH	1	2	2	4	K ² =0.44 F.P=0.604
CERVICAL/VAGINAL TEAR	1	2	2	4	K ² =0.44 F.P=0.604
NONE	48	96	46	92	

12. POSTPARTUM COMPLICATIONS (CASAERAN DELIVERY) AND ITS RELATION WITH BMI

Wound infection was absent in the BMI <30 category and 6% in BMI >30 category. The difference was statistically insignificant with a p value of 0.24.

Hospital stay was prolonged in these 6% patients in BMI >30 category with Post LSCS wound infection. P value was calculated at 0.24 making it statistically insignificant.

TABLE NO. 12 POSTPARTUM COMPLICATIONS (CASAERAN DELIVERY) AND ITS RELATION WITH BMI

COMPLICATION	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE D.F=1
POST LSCS WOUND INFECTION	0	0	3	6	F.P=0.24 K ² =3.09
NONE	50	100	47	94	

TABLE NO. 13 PREVALANCE OF ANEMIA AND ITS RELATION WITH BMI

Prevalence of anemia in BMI < 30 group was 22%, and in the BMI > 30 group was 16%. The difference was statistically insignificant with a p value of 0.444.

TABLE NO. 13 PREVALANCE OF ANEMIA AND ITS RELATION WITH BMI

HB	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE D.F=1
<10	11	22	8	16	K ² =0.58, P=0.444
>10	39	78	42	84	

13.USG ABNORMALITIES AND ITS RELATION WITH BMI

Oligohydramnios in the BMI <30 category was 6%, and in the BMI >30 category was 4%.

Doppler abnormalities in the BMI >30 category was 8%, and these were conspicuously absent in the BMI <30 category. Low lying placenta was found equally in both the groups.

Both oligohydramnios and Doppler changes were seen in 4% patients in BMI<30 group and in 2% women in BMI>30 group.

The difference was statistically insignificant with a p value of 0.56.

TABLE NO. 13 USG ABNORMALITIES AND ITS RELATION WITH BMI

USG FINDINGS	BMI < 30 no.	BMI < 30 %	BMI > 30 no.	BMI > 30 %	K ² AND P VALUE
NORMAL	44	88	42	84	K ² =0.33, P=0.56
OLIGOHYDRAMNIOS	3	6	2	4	
DOPPLER ABNORMALITY	0	0	4	8	
LOW LYING PLACENTA	1	2	1	2	
OLIGOHYDRAMNIOS AND DOPPLER	2	4	1	2	

DISCUSSION

The body mass index (BMI), or Quetelet index, is used to assess the degree of obesity in a patient, based on an individual's weight and height. It was devised between 1830 and 1850, and is defined as the individual's body weight (in kilograms) divided by the square of his or her height (in meters). The formulae universally used in medicine produces a unit of measure of

kg/m². Because BMI is derived from simple measurements of height and weight, it is clearly inexpensive.

In the recent times, obesity has emerged as a health hazard as excess bodyweight is a major cause of diseases worldwide and increased level of obesity may result in a decline in life expectancy in the future. Some investigators have suggested that certain ethnic groups like Asians may be at risk for comorbidities due to obesity at lower BMI thresholds than for other ethnic groups.

A total of 100 cases, 50 with BMI>30 and 50 with BMI<30 were included in this study undertaken at Kasturba Hospital, Delhi, from April 2011 to April 2012. The primigravidas who presented in the labour room after 28 weeks of gestation were included. The antenatal, intrapartum, postpartum and neonatal assessment was done and outcome of each pregnancy in terms of maternal and perinatal morbidity and mortality were studied.

AGE

In our study, 48% of the BMI > 30 category women were >26 years of age, whereas only 28% of the BMI < 30 group were in the >26 years category. The p value comes out to be 0.039 making the difference statistically significant. Mean age was 25.92 in the BMI>30 group compared with 24.2 in the BMI<30 group. This could be due to the age related weight gain in these patients.

Our results were comparable with Meher-Un-Nisa et al (2009) who reported that average age of obese patients was 25.2 and that of non obese was 24.1, showing that obesity was more often found in women of higher age.²⁹

ANTEPARTUM COMPLICATIONS

Preeclampsia

In our study, the frequency of preeclampsia remained significantly high in BMI > 30 category as compared to BMI < 30 group. The frequency of preeclampsia was 38% in the BMI > 30 category and 8% in the BMI < 30 category. The difference was statistically significant with a p value of 0.0003. Eclampsia was found in 2% patients in the BMI >30 category, and was not found in BMI<30 category. P value of 1 was statistically insignificant.

Our results were comparable with Voigt et al (2008) who found that 37.9% patients in the BMI>30 category had preeclampsia and 1.2% in the BMI < 25 category had preeclampsia.²⁶

Ehrental DB (2011) also concluded that preeclampsia was more common in the obese with a p value of less than 0.0001.³⁸

Also, Baeten JM et al (2001) found that incidence of eclampsia increased with increasing BMI.²¹

Retinopathy

Retinopathy was 6% in the BMI >30 category, and 2% in BMI <30. The difference was statistically insignificant with a p value of 0.617.

This could be because of the higher prevalence of preeclampsia and GDM in the BMI>30 group as these are associated with retinal changes.

GDM

Results of our study show that rate of gestational diabetes mellitus in women with BMI>30 was 6% whereas it was only 2% in the BMI<30 category. The difference however was insignificant with a with a p value of 0.617.

Our results were similar to the study by Bianco AT et al (1998) reported in their study of 613 obese patients, a higher prevalence of gestational diabetes mellitus in the obese group (14.2%) as compared to their non obese group (1.2%).⁹²

Kongubol A and Phupong V (2011) said that prepregnancy obesity without metabolic problems did not increase the risk for GDM.⁴¹

The risk of Diabetes Mellitus increases as the age increases, especially after 45 years of age. As our study group was of a younger age group, rates of diabetes were much lower.

IUGR

In our study, the frequency of IUGR remained insignificantly high in BMI > 30 category at 6%, compared to 4% in BMI < 30 group. The difference was statistically insignificant with a p value of 0.646.

This could be due to a possible confounding effect of preeclampsia, as obese patients have higher prevalence of preeclampsia, which has been associated with IUGR for long.

Our results corroborated with the findings of Perlow JH (1992) who reported intrauterine growth retardation at 8.1% in the obese compared to 0.9% in the non obese. However, when those massively obese pregnant women with diabetes and/or hypertension antedating pregnancy are excluded from analysis, no statistically significant differences in perinatal outcome persisted.⁷²

Also, Baeten JM et al (2001) who found that IUGR in the overweight and obese group was 5.1% and 5.6% respectively, compared with 6.1% in the non obese group.²¹

Preterm labour pains

Preterm labour pains occurred in 6% of the pregnancies with BMI < 30 and 10% in the BMI >30 category. The difference was statistically insignificant with a p value of 0.7149.

Our study was similar to a study by Aly H et al (2010) who reported that mothers with obesity and morbid obesity were more likely to deliver prematurely (16.7 and 20.3%, respectively) when compared with non obese women (14.5%). However, when controlling for confounders, obesity and morbid obesity were not associated with prematurity.⁸¹

Similar results were reported by Mandal D et al (2011) who said that preterm labor in less than 34 week gestation was more common in the obese patients.⁹³

MALPRESENTATIONS

Our study reported 4% patients with malpresentation in the BMI>30 group and 2% in the the BMI<30 group. The difference was statistically insignificant with a p value of 1. There was a single patient with breech presentation in the BMI<30 group and 2 patients with malpresentations in the BMI>30 group (1 breech and 1 transverse lie).

Our results were similar to those of Sheiner E etal (2004) reported malpresentations at a significantly higher rate in the obese gravida ($P < 0.001$).⁹⁴

PERIOD OF GESTATION

Preterm labour pains occurred in 6% of the pregnancies with BMI < 30 and 10% in the BMI >30 category. The difference was statistically insignificant with a p value of 0.7149.

Mothers reaching beyond term (post term) were 4% in the BMI<30 group and no posterm patients were seen in the BMI>30 group. The difference was statistically insignificant with a p value of 0.4949.

Our results were inconsistent with those of Caughey AB etal (2009) who reported gestation beyond 41 weeks to include obesity as a cause(adjusted odds ratio [aOR], 1.26; 95% confidence interval [CI], 1.16-1.37). This could be due to the possible confounding effect of preeclampsia which led to earlier inductions/LSCS in the BMI>30 women.⁹⁵

INTRAPARTUM COMPLICATIONS

Fetal distress

Fetal distress was present in 6% patients with BMI <30 category and was absent in the BMI>30 group. The difference was statistically insignificant with a p value of 0.24.

In contrast, Bianco AT etal (1998) found increased incidence of fetal distress(12.4%) in the obese as compared to non obese (8.7%).⁹²

This difference could be due to earlier detection of compromised fetus in the BMI>30 category, due to higher degree of clinical suspicion in view of associated complications like preeclampsia, IUGR. Earlier inductions/ elective LSCS in the same could deter any fetal distress from arising in a stressed fetus also. Also, out of the 3 detected fetal distress patients in BMI<30 group, 2 were unbooked patients, not receiving any previous medical care.

NPOL

NPOL was present in 2% patients with BMI <30 category and was absent in the BMI >30. The difference was statistically insignificant with a p value of 1.

Our results were in contrast with those of Bianco AT et al (1998) reported a higher incidence of NPOL (12.9%) in obese as compared to 7.3% in the non obese.⁹²

We actively manage labour patients in our hospital, and any abnormality in progress of labour is quickly detected. The difference in values could be due to the smaller sample size in our study.

Failure of induction

Failure of induction occurred in 2% patients with BMI <30 category and 2% in BMI>30. No statistical analysis could be done due to similar values and it was found at equal frequency in both the groups.

Shoulder dystocia

Shoulder dystocia was present in only 2% of the patients in the BMI>30 category, whereas it was absent in patients with BMI<30. The difference was statistically insignificant with a p value of 1.

Our results were similar to Meher-Un-Nisa et al (2009), who, in their study reported the frequency of shoulder dystocia to be high in overweight, obese and morbidly obese females (1–7%) as compared to normal weight group (0%).²⁹

MODE OF DELIVERY

Results of our study show significantly higher rates of cesarean section in BMI>30 group as compared to those with BMI<30 group (56% versus 22%).

Our results could be compared with those of Pevzner L etal (2009) said that the incidence of cesarean delivery increased from 21.3% in the BMI less than 30 group to 29.8% in the BMI 30-39.9 group and 36.5% in the BMI 40 or higher group. ⁵¹

Also, Kominiarek MA etal (2010) said that the risk for cesarean increased as BMI increased for all subgroups, $P < .001$. The risk for cesarean increased by 5%, 2%, and 5% for nulliparas and multiparas with and without a prior cesarean, respectively, for each 1-kg/m² increase in BMI. ⁵⁴

ANAESTHETIC COMPLICATIONS

Anaesthetic complications occurred in none of the patients in BMI <30 category and in 10.17% of patients with BMI>30. These complications included failure of spinal anaesthesia in 2 patients and need for general anaesthesia in them. Also, 1 patient in BMI>30 category had intraoperative changes in the ECG suggestive of myocardial infarction and was treated for the same. Statistical analysis revealed that p value was 0.545 making the difference insignificant.

Our results matched with Mace HS etal (2011) who found obese pregnant women appear to have increased morbidity and mortality associated with caesarean delivery and general anaesthesia for caesarean delivery in particular, and more anaesthesia-related complications. ⁵⁷

ELECTIVE AND EMERGENCY LSCS

In the BMI<30 group, 27.27% patients had an elective LSCS whereas 72.72% had an emergency LSCS. In BMI>30 group, 35.714% patients had an elective LSCS whereas 64.285% had an emergency LSCS. The results were statistically insignificant with a p value of 0.719.

Our results were inconsistent with that of Bhattacharya etal (2007), who reported 41.5% emergency LSCS in the normal and 58.8% in the obese group. ⁹⁶

Our results were comparable with Elíasdóttir ÓJ et al (2010) who said that obese women have a significantly increased risk of induction of labour and being delivered by cesarean section, both emergent and elective compared to mothers of normal weight and overweight.³⁶

This was because many of the high risk patients with preeclampsia/ IUGR were taken up for elective LSCS directly in our hospital.

Most common reason for casaerean sections in BMI>30 group was preeclampsia with/without IUGR/Doppler abnormalities. Most common reason for casaerean sections in BMI<30 group was Meconium stained liquor intrapartum.

INTRAOPERATIVE LSCS COMPLICATIONS

Intraoperative lower segment caesarean sections were complicated in 2% patients with BMI <30 category and 10% in BMI>30 category. Statistical analysis showed p value of 0.204 making it statistically insignificant. These included difficulty in opening up the patient for LSCS in 4 patients with BMI> 30 and rent in broad ligament in 1 of them. In 1 unbooked patient with BMI<30, we did an emergency LSCS in view of obstructed labour and she had bladder injury intraoperatively.

Our results were similar to those of Perlow JH et al (1994) who reported that massively obese pregnant women undergoing cesarean section were at significantly increased risk for peroperative morbidity.⁷²

Norman JE and Reynolds RM (2011) also found that obesity complicates operative delivery; it makes operative delivery more difficult, increases complications and paradoxically increases the need for operative delivery.⁶⁹

NEED FOR INDUCTION

Inductions were done in 12% of the BMI <30 category and 14% of the the BMI >30 category. The difference was statistically insignificant with a p value of 0.766.

The most common indication for induction in the BMI>30 group was preeclampsia whereas in BMI<30 group was postdatism.

Our results were comparable with Jensen DM et al (2003) reported that the risk of induction of labor was significantly increased in both overweight women (body mass index [BMI] 25.0-29.9 kg/m²) and obese women (BMI \geq 30.0 kg/m²) compared with women who were of normal weight (BMI 18.5-24.9 kg/m²).²²

Also, Elíasdóttir ÓJ et al (2010) who reported that obese women have a significantly increased risk of requiring induction of labour compared with normal weight women.³⁶

POSTPARTUM COMPLICATIONS

PPH

PPH occurred in 2% of the patients with BMI <30 category and in 4% of the patients in the BMI >30 category. The difference was statistically insignificant with a p value of 0.604.

Our results were consistent with those of T.S. Usha Kiran, S. Hemmadi, J. Bethel, J. Evans (2005) who reported an increased risk [quoted as odds ratio (OR) and confidence intervals CI] of maternal complications such as blood loss of more than 500 ml, amounting to postpartum haemorrhage.⁴⁸

Cervical/ Paravaginal tears

Cervical/ Paravaginal tears were present in 2% of the BMI <30 category and 4% in BMI >30 category. The difference was statistically insignificant with a p value of 0.604.

Our results were comparable with Liu X et al (2011) who found a significant increase in postpartum hemorrhage and perineal rupture in obese patients.⁶⁶

WOUND INFECTION

Wound infection was absent in the BMI <30 category and 6% in BMI >30 category. The difference was statistically insignificant with a p value of 0.24.

The local changes, such as an increase in adipose tissue, an increase in local tissue trauma related to retraction, the immune dysfunction, increased association of diabetes with obesity and a

lengthened operative time, may contribute to the increased incidence of surgical site infections caused by obesity.

Our results can be compared with those of Satpathy HK et al (2008) who reported that following Cesarean section delivery, obese women have a higher incidence of wound infection and disruption.⁶³

Alanis MC et al (2010) reported that women with a body mass index $>$ or $=$ 50 kg/m^2 have a much greater risk for cesarean wound complications than previously reported. Avoidance of subcutaneous drains and increased use of transverse abdominal wall incisions should be considered in massively obese parturients to reduce operative morbidity.⁵³

Mandal D et al (2011) said that obese pregnant women were at increased risk of postpartum infection morbidities.⁹³

ANEMIA

Prevalence of anemia in BMI $>$ 30 was 16% and 22% in the BMI $<$ 30 category. The difference was statistically insignificant with a p value of 0.444.

These results could be due to possible nutritional etiology of anemia in the population with BMI $<$ 30.

Our results could be compared with Galtier-Dereure F et al (2000) who reported that anemia appears to occur less often in severely obese pregnant women than in normal-weight pregnant women.⁷³

Aly H et al (2010) found that mothers with obesity and morbid obesity were more likely to have anemia than normal weight women.⁸¹

USG

Oligohydramnios in the BMI $<$ 30 category was 6%, and in the BMI $>$ 30 category was 4%.

Doppler abnormalities in the BMI $>$ 30 category was 8%, and these were conspicuously absent in the BMI $<$ 30 category.

The difference in ultrasound findings remained statistically insignificant with a p value of 0.56.

CONCLUSION

From our study we may conclude that there is a higher prevalence of complications to both the fetus and the mother when BMI is more than 30 in the mother. Women with BMI > 30 had significantly higher age than women with BMI less than 30, and were associated with significantly increased incidence of preeclampsia, caesarean sections, and lower APGAR. There was an insignificant increase in eclampsia, retinopathy, gestational diabetes mellitus, intra uterine growth restriction, preterm labour pains, malpresentations, shoulder dystocia. Also, anaesthetic complications, elective caesarean sections and intraoperative complications, inductions, postpartum hemorrhage, cervical/paravaginal tears, post-operative wound infection, Doppler abnormalities and macrosomia were insignificantly higher in the BMI more than 30 group. The incidence of failed induction and intra uterine deaths was similar in both the groups. The following were insignificantly higher in the BMI less than 30 group: postdatism, fetal distress, non progress of labour, anemia, oligohydramnios, low birth weight, meconium aspiration syndrome, NICU admissions.

Therefore, it is a must for all pregnant and non pregnant women to be aware of the fetomaternal complications arising due to higher Body Mass Index. With proper management of pregnant women with a higher BMI, improvement in awareness amongst the women and increasing their accessibility to medical facilities, maternal and perinatal morbidity and mortality can be minimized. Preconceptional weight loss and limited pregnancy weight gain can be helpful in achieving the goal we all strive for, a healthy mother and a healthy baby.

ACKNOWLEDGEMENT

I had the proud privilege and honor of having worked under the excellent guidance and supervision of my esteemed teacher and mentor Dr Asha Aggarwal, Head of Department, Department of Obstetrics and Gynaecology, Kasturba Hospital for her valuable advice, inspiring guidance, stimulating suggestions and positive criticisms, motivational approach and meticulous supervision at each step of this work.

I would also like to extend my sincere thanks to Dr Gurcharan Kaur, Senior Specialist and Head of Unit, for her guidance, painstaking supervision and kind help. I am really thankful for providing comprehensive knowledge and stimulating new thoughts on the subject, which was helpful in completing this work.

I would like to thank Dr. Khan Amir Maroof, who was a constant source of inspiration and encouragement. Also, I would like to thank all my colleagues and friends for their valuable support and cooperation. I am also grateful for the technical, administrative and other non-teaching staff for their cooperation and support in my endeavors.

Last but not the least; I shall ever remain indebted to my mother Mrs. Anjla Lakhanpal and my father Mr. Raman Lakhanpal, and my little sister Miss. Nupur Lakhanpal for their untiring and valuable support, inspirations and encouragement in completing this work.

Above all, I record my utmost gratitude to all the patients for their voluntary cooperation and active participation in the study.

Dr Shuchi Lakhanpal

INDEX OF REFERENCES

1. James WP. WHO recognition of the global obesity epidemic. *Int J Obes (Lond)* 2008;32 Suppl 7:S120-6.
2. Haslam DW, James WP. Obesity. *Lancet* 2005;366:1197–1209.

3. Eknoyan, Garabed. Adolphe Quetelet (1796–1874)—the average man and indices of obesity. *Nephrol Dial Transplant* 2008;23(1):47–51.
4. Fidanza F, Karvonen MJ, Kimura N, Taylor HL. Indices of relative weight and obesity. *J Chronic Dis* 1972;25(6): 329–43.
5. Bloomgarden Z. Prevention of Obesity and Diabetes. *Diabetes Care* 2003;26 (11): 3172–3178.
6. Haslam DW, James WP. Obesity. *Lancet* 2005;366 (9492): 1197–209.
7. Mazzone T, Fantuzzi G. *Adipose Tissue And Adipokines in Health And Disease (Nutrition and Health)*. Totowa, NJ: Humana Press 2006: 222.
8. Haslam D. Obesity: a medical history. *Obes Rev* 2007;8 Suppl 1: 31–6.
9. Hill HG, Smith JF. The Relation of Fatness to Sterility. *BJOG* 1930;37(2): 256-71.
10. Gill AM. Unclassified Types of Obesity. *Acta Medica Scandinavica* 1938;95(S90) 257–68.
11. Bray GA. Complications of Obesity. *Annals of Internal Medicine* 1985;103:1052-62.
12. Brown PJ, Konner M. An Anthropological Perspective on Obesity. *Human Obesity* 1987;499: 29–46.
13. Pagano R, La Vecchia C. Overweight and obesity in Italy, 1990-91. *Int J Obes Relat Metab Disord* 1994;18(10):665-9.
14. Gidding SS, Leibel RL, Daniels S, Rosenbaum M, Horn LV, Marx GR. Understanding Obesity in Youth. *Circulation* 1996;94:3383-87.
15. Bouchard C. Genetics of Human Obesity: Recent Results from Linkage Studies. *J Nutr* 1997; 127 (9): 1887S-1890S.
16. Sturm R. Increases in morbid obesity in the USA: 2000–2005. *Public Health* 2007;121(7): 492–6.
17. Afridi AK, Safari M, Chattel MMAK, Khan A. Health Risk of Overweight and Obesity- An Overview. *Pakistan Journal of Nutrition* 2003;2(6):350-360

18. Jarvie E, Hauguel-de-Mouzon S, Nelson SM, Sattar N, Catalano PM, Freeman DJ. Lipotoxicity in obese pregnancy and its potential role in adverse pregnancy outcome and obesity in the offspring. *Clin Sci (Lond)* 2010;119(3):123-9.
19. Dohrmann KR, Sally, Lederman A. Weight Gain in Pregnancy. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*. 1986;15(6):446–53.
20. Ekblad U, Grenman S. Maternal weight, weight gain during pregnancy and pregnancy outcome. *Int J Gynaecol Obstet* 1992;39(4):277-83.
21. Baeten JM, Bukusi EA, Lambe M. Pregnancy complications and outcomes among overweight and obese nulliparous women. *Am J Public Health* 2001; 91(3): 436–40.
22. Jensen DM, Ovesen P, Beck-Nielsen H, Mølsted-Pedersen L, Sørensen B, Vinter C et al. Gestational Weight Gain and Pregnancy Outcomes in 481 Obese Glucose-Tolerant Women. *Diabetes Care*. 2005;28(9)2118-2122.
23. Bodnar LM, Catov JM, Roberts JM, Simhan HN. Prepregnancy Obesity Predicts Poor Vitamin D Status in Mothers and Their Neonates. *J Nutr* 2007;137(11):2437-2442.
24. Burstein E, Levy A, Mazor M, Wiznitzer A, Sheiner E. Pregnancy outcome among obese women: a prospective study. *Am J Perinatol*. 2008;25(9):561-6.
25. Leung TY. Trends in maternal obesity and associated risks of adverse pregnancy outcomes in a population of Chinese women. *BJOG* 2008;115(12):1529-37.
26. Voigt M, Straube S, Zygmunt M, Krafczyk B, Schneider KT, Briese V. Obesity and pregnancy--a risk profile. *Z Geburtshilfe Neonatol*. 2008;212(6):201-5.
27. Yogev Y, Visser GH. Obesity, gestational diabetes and pregnancy outcome. *Semin Fetal Neonatal Med* 2009;14(2):77-84.

28. Joy S, Istwan N, Rhea D, Desch C, Stanziano G. The impact of maternal obesity on the incidence of adverse pregnancy outcomes in high-risk term pregnancies. *Am J Perinatol* 2009;26(5):345-9.
29. Meher-Un-Nisa, Aslam M, Ahmed SR, Rajab M, Kattea L. Impact of obesity on fetomaternal outcome in pregnant saudi females. *Int J Health Sci (Qassim)* 2009;3(2):187-95.
30. Khashan AS, Kenny LC. The effects of maternal body mass index on pregnancy outcome. *Eur J Epidemiol* 2009;24(11):697-705.
31. Ulman-Włodarz I, Nowosielski K, Romanik M, Pozowski J, Krawczyk P. Pregnancy and delivery course in overweight and obese women. *Ginekol Pol* 2009;80(10):744-51.
32. Briese V, Voigt M, Hermanussen M, Wittwer-Backofen U. Morbid obesity: pregnancy risks, birth risks and status of the newborn. *Homo* 2010;61(1):64-72.
33. Kanadys WM, Leszczyńska-Gorzela B, Oleszczuk J. Obesity among women. Pregnancy after bariatric surgery: a qualitative review. *Ginekol Pol* 2010;81(3):215-23.
34. Nelson SM, Matthews P, Poston L. Maternal metabolism and obesity: modifiable determinants of pregnancy outcome. *Hum Reprod Update* 2010;16(3):255-75.
35. Lapolla A, Marangon M, Dalfrà MG, Segato G, De Luca M, Fedele D et al. Pregnancy outcome in morbidly obese women before and after laparoscopic gastric banding. *Obes Surg* 2010;20(9):1251-7.
36. Elíasdóttir ÓJ, Harðardóttir H, Þórkelsson Þ. The effect of maternal weight on pregnancy outcome. *Laeknabladid* 2010;96(11):691-6.
37. Choi SK, Park IY, Shin JC. The effects of pre-pregnancy body mass index and gestational weight gain on perinatal outcomes in Korean women: a retrospective cohort study. *Reprod Biol Endocrinol* 2011;18(9):6.
38. Ehrenthal DB, Jurkovitz C, Hoffman M, Jiang X, Weintraub WS. Prepregnancy body mass index as an independent risk factor for pregnancy-induced hypertension. *J Womens Health (Larchmt)* 2011;20(1):67-72.

39. Misra VK, Trudeau S. The influence of overweight and obesity on longitudinal trends in maternal serum leptin levels during pregnancy. *Obesity (Silver Spring)* 2011;19(2):416-21.
40. Simmons D. Diabetes and obesity in pregnancy. *Best Pract Res Clin Obstet Gynaecol* 2011;25(1):25-36.
41. Kongubol A, Phupong V. Prepregnancy obesity and the risk of gestational diabetes mellitus. *BMC Pregnancy Childbirth* 2011;11:59.
42. Gaillard R, Steegers EA, Hofman A, Jaddoe VW. Associations of maternal obesity with blood pressure and the risks of gestational hypertensive disorders. The Generation R Study. *J Hypertens* 2011;29(5):937-44.
43. Olivarez SA, Ferres M, Antony K, Mattewal A, Maheshwari B, Sangi-Haghpeykar H et al Obstructive sleep apnea screening in pregnancy, perinatal outcomes, and impact of maternal obesity. *Am J Perinatol* 2011;28(8):651-8.
44. Biering K, Nøhr EA, Olsen J, Andersen AM, Hjøllund NH, Juhl M. Pregnancy-related pelvic pain is more frequent in women with increased body mass index. *Acta Obstet Gynecol Scand.* 2011;90(10):1132-9.
45. Wilson AT, Reilly CS. Anaesthesia and the obese patient. *Int J Obes Relat Metab Disord.* 1993;17(8):427-35.
46. Cohen S, Gondret R, Mankikian B. Massive obesity and anesthesia. *Rev Prat.* 1993;43(15):1950-5.
47. Gautam PL, Kathuria S, Kaul TK. *Acta Anaesthesiol Scand.* Infiltration block for caesarean section in a morbidly obese parturient 1999;43(5):580-1.
48. Usha Kiran TS, Hemmadi S , Bethel J, Evans J. Outcome of pregnancy in a woman with an increased body mass index 2005;112(6):768-772.
49. Haeri S, Guichard I, Baker AM, Saddlemire S, Boggess KA. The effect of teenage maternal obesity on perinatal outcomes. *Obstet Gynecol* 2009;113(2 Pt 1):300-4.

50. Magriples U, Kershaw TS, Rising SS, Westdahl C, Ickovics JR. The effects of obesity and weight gain in young women on obstetric outcomes. *Am J Perinatol* 2009;26(5):365-71.
51. Pevzner L, Powers BL, Rayburn WF, Rumney P, Wing DA. Effects of maternal obesity on duration and outcomes of prostaglandin cervical ripening and labor induction. *Obstet Gynecol.* 2009;114(6):1315-21.
52. Robinson CJ, Hill EG, Alanis MC, Chang EY, Johnson DD, Almeida JS. Examining the effect of maternal obesity on outcome of labor induction in patients with preeclampsia. *Hypertens Pregnancy* 2010;29(4):446-56.
53. Alanis MC, Villers MS, Law TL, Steadman EM, Robinson CJ. Complications of cesarean delivery in the massively obese parturient. *Am J Obstet Gynecol* 2010;203(3):271.e1-7.
54. Kominiarek MA, Vanveldhuisen P, Hibbard J, Landy H, Haberman S, Learman L et al. The maternal body mass index: a strong association with delivery route. *Am J of Obs & Gyne* 2010;203(3):264.e1-264.e7.
55. Arrowsmith S, Wray S, Quenby S. Maternal obesity and labour complications following induction of labour in prolonged pregnancy. *BJOG* 2011;118(5):578-88.
56. Beyer DA, Amari F, Lüdders DW, Diedrich K, Weichert J. Obesity decreases the chance to deliver spontaneously. *J Arch Gynecol Obstet* 2011;283(5):981-8.
57. Mace HS, Paech MJ, McDonnell NJ. Obesity and obstetric anaesthesia. *Anaesth Intensive Care* 2011;39(4):559-70.
58. Alexandra P, Vassilios B, Alexandra V, George K, Vassiliki L, Chryssa B. Population-based trends of pregnancy outcome in obese mothers: what has changed over 15 years. *Obesity (Silver Spring)* 2011;19(9):1861-5.
59. Vricella LK, Louis JM, Mercer BM, Bolden N. Impact of morbid obesity on epidural anesthesia complications in labor. *Am J Obstet Gynecol* 2011;205(4):370.e1-6.
60. Porat S, Maxwell C, Sermer M, Farine D. The degree of obesity effect on labor outcome following induction. *Am J of Obs & Gyne* 2012;206(1)Suppl:S84-S85.

61. Sebastián Manzanares G, Angel Santalla H, Irene Vico Z, López Criado MS, Alicia Pineda L, José Luis Gallo V. Abnormal maternal body mass index and obstetric and neonatal outcome. *J Matern Fetal Neonatal Med* 2012;25(3):308-12.
62. Lacoursiere DY, Hutton A, Varner M. The association of obesity, body image and postpartum depression. *Am J of Obs & Gyne* 2007;197(6)Suppl:S93.
63. Satpathy HK, Fleming A, Frey D, Barsoom M, Satpathy C, Khandalavala J. Maternal obesity and pregnancy. *Postgrad Med* 2008;120(3):E01-9.
64. Nohr EA, Timpson NJ, Andersen CS, Davey Smith G, Olsen J, Sørensen TI. Severe obesity in young women and reproductive health: the Danish National Birth Cohort. *PLoS One* 2009;4(12):e8444.
65. Baron CM, Girling LG, Mathieson AL, Menticoglou SM, Seshia MM, Cheang MS et al. Obstetrical and neonatal outcomes in obese parturients. *J Matern Fetal Neonatal Med* 2010;23(8):906-13.
66. Liu X, Du J, Wang G, Chen Z, Wang W, Xi Q. Effect of pre-pregnancy body mass index on adverse pregnancy outcome in north of China. *Arch Gynecol Obstet.* 2011;283(1):65-70.
67. Novaes JF, Lamounier JA, Colosimo EA, Franceschini SC, Priore SE. Breastfeeding and obesity in Brazilian children. *Eur J Public Health* 2011; (Epub ahead of print)
68. Regnault N, Botton J, Blanc L, Hankard R, Forhan A, Goua V. Determinants of neonatal weight loss in term-infants: specific association with pre-pregnancy maternal body mass index and infant feeding mode. *Arch Dis Child Fetal Neonatal* 2011;96(3):F217-22.
69. Norman JE, Reynolds RM. The consequences of obesity and excess weight gain in pregnancy. *Proc Nutr Soc* 2011;70(4):450-6.
70. Taper LJ, Hayes M, Rogers CS, Fray RB. Influence of Maternal Weight, Smoking, and Socioeconomic Status on Infant Triceps Skinfold Thickness and Growth During the First Year. *Birth* 1984;11(2):97-101.

71. Rössner S, Öhlin A. Maternal Body Weight And Relation To Birth Weight. *Acta Obstetricia et Gynecologica Scandinavica* 1990;69(6):475–478.
72. Perlow JH, Morgan MA, Montgomery D, Towers CV, Porto M. Perinatal outcome in pregnancy complicated by massive obesity. *Am J Obstet Gynecol* 1992;167(4 Pt 1):958-62.
73. Galtier-Dereure F, Boegner C, Jacques Bringer. Obesity and pregnancy: complications and cost. *American Journal of Clinical Nutrition* 2000;71(5):1242S-1248s.
74. A.S. Kumari. Pregnancy outcome in women with morbid obesity. *International Journal of Gynecology & Obstetrics* 2001;73(2):101-107.
75. Bringer J, Galtier F, Raingard I, Boulot P, Renard E. Pregnancy and overweight: underestimated consequences? *Bull Acad Natl Med* 2008;192(4):673-87.
76. Oken E. Excess gestational weight gain amplifies risks among obese mothers. *Epidemiology* 2009;20(1):82-3.
77. Wax JR. Risks and management of obesity in pregnancy: current controversies. *Curr Opin Obstet Gynecol* 2009;21(2):117-23.
78. Blomberg MI, Källén B. Maternal obesity and morbid obesity: the risk for birth defects in the offspring. *Birth Defects Res A Clin Mol Teratol* 2010;88(1):35-40.
79. Chen M, McNiff C, Madan J, Goodman E, Davis JM, Dammann O. Maternal obesity and neonatal Apgar scores. *J Matern Fetal Neonatal Med* 2010;23(1):89-95.
80. Wise LA, Palmer JR, Heffner LJ, Rosenberg L. Prepregnancy body size, gestational weight gain, and risk of preterm birth in African-American women. *Epidemiology* 2010 Mar;21(2):243-52.
81. Aly H, Hammad T, Nada A, Mohamed M, Bathgate S, El-Mohandes A. Maternal obesity, associated complications and risk of prematurity. *J Perinatol*. 2010 Jul;30(7):447-51.
82. Alanis MC, Goodnight WH, Hill EG, Robinson CJ, Villers MS, Johnson DD. Maternal super-obesity (body mass index \geq 50) and adverse pregnancy outcomes. *Acta Obstet Gynecol Scand* 2010;89(7):924-30.

83. McDonald SD, Han Z, Mulla S, Beyene J; Knowledge Synthesis Group. Overweight and obesity in mothers and risk of preterm birth and low birth weight infants: systematic review and meta-analyses. *BMJ* 2010;341:c3428.
84. Narchi H, Skinner A. Overweight and obesity in pregnancy do not adversely affect neonatal outcomes: new evidence. *J Obstet Gynaecol.* 2010;30(7):679-86.
85. Lapolla A, Bonomo M, Dalfrà MG, Parretti E, Mannino D, Mello G. Prepregnancy BMI influences maternal and fetal outcomes in women with isolated gestational hyperglycaemia: a multicentre study. *Diabetes Metab* 2010;36(4):265-70.
86. Kitsantas P, Pawloski LR, Gaffney KF. Maternal prepregnancy body mass index in relation to Hispanic preschooler overweight/obesity. *Eur J Pediatr* 2010;169(11):1361-8.
87. Begum KS, Sachchithanatham K, De Somsubhra S. Maternal obesity and pregnancy outcome. *Clin Exp Obstet Gynecol* 2011;38(1):14-20.
88. Stacey T, Thompson JM, Mitchell EA, Ekeroma AJ, Zuccollo JM, McCowan LM. Relationship between obesity, ethnicity and risk of late stillbirth: a case control study. *BMC Pregnancy Childbirth* 2011;11:3.
89. Roman AS, Rebarber A, Fox NS, Klauser CK, Istwan N, Rhea D et al. The effect of maternal obesity on pregnancy outcomes in women with gestational diabetes. *J Matern Fetal Neonatal Med* 2011;24(5):723-7.
90. Modi N, Murgasova D, Ruager-Martin R, Thomas EL, Hyde MJ, Gale C et al. The influence of maternal body mass index on infant adiposity and hepatic lipid content. *Pediatr Res* 2011;70(3):287-91.
91. Yessoufou A, Moutairou K. *Exp Diabetes Res.* Maternal diabetes in pregnancy: early and long-term outcomes on the offspring and the concept of "metabolic memory" 2011;2011:218598.

92. Bianco AT, Smilen SW, Davis Y, Lopez S, Lapinski R, Lockwood CJ. Pregnancy outcome and weight gain recommendations for the morbidly obese woman. Obstet Gynecol 1998;91(1):97-102.

93. Mandal D, Manda S, Rakshi A, Dey RP, Biswas SC, Banerjee A. Maternal obesity and pregnancy outcome: a prospective analysis. J Assoc Physicians India 2011;59:486-9.

94. Sheiner E, Levy A, Menes TS, Silverberg D, Katz M, Mazor M. Maternal obesity as an independent risk factor for caesarean delivery. Paediatr Perinat Epidemiol 2004;18(3):196-201.

95. Caughey AB, Stotland NE, Washington AE, Escobar GJ. Who is at risk for prolonged and postterm pregnancy? Am J Obstet Gynecol 2009;200(6):683.e1-5.

96. Bhattacharya S, Campbell DM, Liston WA, Bhattacharya S. Effect of Body Mass Index on pregnancy outcomes in nulliparous women delivering singleton babies. BMC Public Health 2007; 7:168.

